

Strong Interactive Massive Particles from a Strong Coupled Theory

Maxim Yu. KHLOPOV* and Chris KOUVARIS†

**Center for Cosmoparticle physics “Cosmion”, 125047, Moscow, Russia
Moscow Engineering Physics Institute, 115409 Moscow, Russia, and
APC laboratory 10, rue Alice Domon et Léonie Duquet 75205 Paris Cedex 13, France,*

*†CERN Theory Division, CH-1211 Geneva 23, Switzerland,
University of Southern Denmark, Campusvej 55, DK-5230 Odense, Denmark and
The Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen, Denmark*

Abstract

Minimal walking technicolor models can provide a nontrivial solution for cosmological dark matter, if the lightest technibaryon is doubly charged. Technibaryon asymmetry generated in the early Universe is related to baryon asymmetry and it is possible to create excess of techniparticles with charge (-2) . These excessive techniparticles are all captured by ${}^4\text{He}$, creating *techni-O-helium* $tOHe$ “atoms”, as soon as ${}^4\text{He}$ is formed in Big Bang Nucleosynthesis. The interaction of techni-O-helium with nuclei opens new paths to the creation of heavy nuclei in Big Bang Nucleosynthesis. Due to the large mass of technibaryons, the $tOHe$ “atomic” gas decouples from the baryonic matter and plays the role of dark matter in large scale structure formation, while structures in small scales are suppressed. Nuclear interactions with matter slow down cosmic techni-O-helium in Earth below the threshold of underground dark matter detectors, thus escaping severe CDMS constraints. On the other hand, these nuclear interactions are not sufficiently strong to exclude this form of Strongly Interactive Massive Particles by constraints from the XQC experiment. Experimental tests of this hypothesis are possible in search for $tOHe$ in balloon-borne experiments (or on the ground) and for its charged techniparticle constituents in cosmic rays and accelerators. The $tOHe$ “atoms” can cause cold nuclear transformations in matter and might form anomalous isotopes, offering possible ways to exclude (or prove?) their existence.

*Electronic address: maxim.khlopov@roma1.infn.it

†Electronic address: kouvaris@nbi.dk

I. INTRODUCTION

The question of the existence of new quarks and leptons is among the most important in the modern high energy physics. This question has an interesting cosmological aspect. If these quarks and/or charged leptons are stable, they should be present around us and the reason for their evanescent nature should be found.

Recently, at least three elementary particle frames for heavy stable charged quarks and leptons were considered: (a) A heavy quark and heavy neutral lepton (neutrino with mass above half the Z -boson mass) of a fourth generation [1, 2], which can avoid experimental constraints [3, 4], and form composite dark matter species [5, 6, 7, 8]; (b) A Glashow's "Sinister" heavy tera-quark U and tera-electron E , which can form a tower of tera-hadronic and tera-atomic bound states with "tera-helium atoms" ($UUUEE$) considered as dominant dark matter [9, 10]; (c) AC-leptons, based on the approach of almost-commutative geometry [11, 12], that can form evanescent AC-atoms, playing the role of dark matter [11, 13, 14].

In all these recent models, the predicted stable charged particles escape experimental discovery, because they are hidden in elusive atoms, composing the dark matter of the modern Universe. It offers a new solution for the physical nature of the cosmological dark matter. Here we show that such a solution is possible in the framework of walking technicolor models [15, 16, 17, 18, 19, 20] and can be realized without an *ad hoc* assumption on charged particle excess, made in the approaches (a)-(c).

This approach differs from the idea of dark matter composed of primordial bound systems of superheavy charged particles and antiparticles, proposed earlier to explain the origin of Ultra High Energy Cosmic Rays (UHECR) [21]. To survive to the present time and to be simultaneously the source of UHECR, superheavy particles should satisfy a set of constraints, which in particular exclude the possibility that they possess gauge charges of the standard model.

The particles considered here, participate in the Standard Model interactions and we show how the problems, related to various dark matter scenarios with composite atom-like systems, can find an elegant solution on the base of the minimal walking technicolor model.

The approaches (b) and (c) try to escape the problems of free charged dark matter particles [22] by hiding opposite-charged particles in atom-like bound systems, which interact weakly with baryonic matter. However, in the case of charge symmetry, when primordial

abundances of particles and antiparticles are equal, annihilation in the early Universe suppresses their concentration. If this primordial abundance still permits these particles and antiparticles to be the dominant dark matter, the explosive nature of such dark matter is ruled out by constraints on the products of annihilation in the modern Universe [3, 13]. Even in the case of charge asymmetry with primordial particle excess, when there is no annihilation in the modern Universe, binding of positive and negative charge particles is never complete and positively charged heavy species should retain. Recombining with ordinary electrons, these heavy positive species give rise to cosmological abundance of anomalous isotopes, exceeding experimental upper limits. To satisfy these upper limits, the anomalous isotope abundance on Earth should be reduced, and the mechanisms for such a reduction are accompanied by effects of energy release which are strongly constrained, in particular, by the data from large volume detectors.

These problems of composite dark matter models [9, 11] revealed in [3, 5, 10, 13], can be avoided, if the excess of only -2 charge A^{--} particles is generated in the early Universe. Here we show that in walking technicolor models, technilepton and technibaryon excess is related to baryon excess and the excess of -2 charged particles can appear naturally for a reasonable choice of model parameters. It distinguishes this case from other composite dark matter models, since in all the previous realizations, starting from [9], such an excess was put by hand to saturate the observed cold dark matter (CDM) density by composite dark matter.

After it is formed in Big Bang Nucleosynthesis, ${}^4\text{He}$ screens the A^{--} charged particles in composite (${}^4\text{He}^{++}A^{--}$) “atoms”. These neutral primordial nuclear interacting objects saturate the modern dark matter density and play the role of a nontrivial form of strongly interacting dark matter [23, 24]. The active influence of this type of dark matter on nuclear transformations seems to be incompatible with the expected dark matter properties. However, it turns out that the considered scenario is not easily ruled out [5, 13] and challenges the experimental search for techni-O-helium and its charged techniparticle constituents.

The structure of the present paper is as follows. Starting with a review of possible dark matter candidates offered by the minimal walking technicolor model, we reveal the possibility for the lightest techniparticle(s) to have electric charge ± 2 (Section II). In Section III we show how the minimal technicolor model can provide substantial excess of techniparticles with electric charge -2 . In Section IV we show how all these -2 charge particles can be captured

by ${}^4\text{He}$, after its formation in the Standard Big Bang Nucleosynthesis (SBBN), making neutral techni-O-helium “atoms” that can account for the modern dark matter density. Techni-O-helium catalyzes a path for heavy element formation in SBBN, but we stipulate in Section IV a set of arguments, by which the considered scenario can avoid immediate contradiction with observations. Gas of heavy techni-O-helium “atoms” decouples from the plasma and radiation only at a temperature about few hundreds eV, so that small scale density fluctuations are suppressed and gravitational instability in this gas develops more close to warm dark matter, rather than to cold dark matter scenario (subsection A of Section V). We further discuss in Section V the possibility to detect charged techniparticle components of cosmic rays (subsection B), effects of techni-O-helium catalyzed processes in Earth (subsection C), and possibilities of direct searches for techni-O-helium (subsection D). The problems, signatures, and possible experimental tests of the techni-O-helium Universe are considered in Section VI. Details of our calculations are presented in the Appendices 1 and 2.

II. DARK MATTER FROM WALKING TECHNICOLOR

The minimal walking technicolor model [15, 16, 17, 18, 19, 20] has two techniquarks, i.e. up U and down D , that transform under the adjoint representation of an $SU(2)$ technicolor gauge group. The global symmetry of the model is an $SU(4)$ that breaks spontaneously to an $SO(4)$. The chiral condensate of the techniquarks breaks the electroweak symmetry. There are nine Goldstone bosons emerging from the symmetry breaking. Three of them are eaten by the W and the Z bosons. The remaining six Goldstone bosons are UU , UD , DD and their corresponding antiparticles. For completeness UU is $U_\alpha^\top C U_\beta \delta^{\alpha\beta}$, where C is the charge conjugate matrix and the Greek indices denote technicolor states. For simplicity in our notation we omit the contraction of Dirac and technicolor indices. Since the techniquarks are in the adjoint representation of the $SU(2)$, there are three technicolor states. The UD and DD have similar Dirac and technicolor structure. The pions and kaons which are the Goldstone bosons in QCD carry no baryon number since they are made of pairs of quark-antiquark. However in our case, the six Goldstone bosons carry technibaryon number since they are made of two techniquarks or two anti-techniquarks. This means that if no processes violate the technibaryon number, the lightest technibaryon will be stable.

The electric charges of UU , UD , and DD are given in general by $y + 1$, y , and $y - 1$ respectively, where y is an arbitrary real number. For any real value of y , gauge anomalies are cancelled [20]. The model requires in addition the existence of a fourth family of leptons, i.e. a “new neutrino” ν' and a “new electron” ζ in order to cancel the Witten global anomaly. Their electric charges are in terms of y respectively $(1 - 3y)/2$ and $(-1 - 3y)/2$. The effective theory of this minimal walking technicolor model has been presented in [19, 25].

There are several possibilities for a dark matter candidate emerging from this minimal walking technicolor model. For the case where $y = 1$, the D techniquark (and therefore also the DD boson) become electrically neutral. If one assumes that DD is the lightest technibaryon, then it is absolutely stable, because there is no way to violate the technibaryon number apart from the sphalerons that freeze out close to the electroweak scale. This scenario was studied in Refs. [19, 20]. It was shown that DD can provide the full dark matter density if its mass is of the order of TeV. The exact value of the mass of DD depends on the temperature where sphalerons freeze out, and on the ratios L/B and L'/B , where L and L' are the lepton number and the lepton number of the fourth lepton family respectively, and B is the baryon number. However, this scenario is ruled out by the CDMS experiment, if DD accounts for 100% of the dark matter density. The reason is that since DD has a Spin Independent (SI) interaction with nuclei, it can scatter coherently in underground dark matter detectors, raising the elastic cross section. Such a cross section is already excluded by CDMS, if we accept that the local dark matter density ranges between $0.2 - 0.4 \text{ GeV}/\text{cm}^3$. However, if DD is a subdominant component, contributing up to 20% of the total dark matter density, it cannot yet be ruled out.

Within the same model and electric charge assignment, there is another possibility. Since both techniquarks and technigluons transform under the adjoint representation of the $SU(2)$ group, it is possible to have bound states between a D and a technigluon G . The object $D^\alpha G^\alpha$ (where α denotes technicolor states) is technicolorless. If such an object has a Majorana mass, then it can account for the whole dark matter density without being excluded by CDMS, due to the fact that Majorana particles have no SI interaction with nuclei and their non-coherent elastic cross section is very low for the current sensitivity of detectors [26]. We should emphasize that nonzero Majorana mass means that the technibaryon number is not protected, as in the previous case. For this scenario to be true, the bound state of DG should be lighter than DD . The lack of tools in order to study the spectrum of the

theory makes hard to decisively conclude if this is true. Lattice calculations have difficulties to study objects like DG and evidently perturbation techniques cannot apply. However, in studies of Super Yang Mills models with supersymmetry softly broken, it has been argued that a Majorana mass for the gluino makes the λG (λ being the gluino) lighter than the $\lambda\lambda$ [27]. If we transfer directly these results in our case, it would mean that DG is lighter than DD as long as D has a Majorana mass. Of course this argument can be only taken as an indication, since the considered walking technicolor model is not supersymmetric and because D in principle has also a Dirac mass. On the other hand, if the Majorana mass is zero, the above argument cannot be applied. In this case it is more natural to expect that DD (or UU and UD), which is a Goldstone boson, is the lightest technibaryon. That might imply that DG is unstable.

Finally, if one choose $y = 1/3$, ν' has zero electric charge. In this case the heavy fourth Majorana neutrino ν' can play the role of a dark matter particle. This scenario was explored first in [28] and later in [26]. It was shown that indeed the fourth heavy neutrino can provide the dark matter density without being excluded by CDMS [29] or any other experiment. This scenario allows the possibility for new signatures of weakly interacting massive particle annihilation [30].

In this paper we study a case that resembles mostly the first one mentioned above, that is $y = 1$ and the Goldstone bosons UU , UD , and DD have electric charges 2, 1, and 0 respectively. In addition for $y = 1$, the electric charges of ν' and ζ are respectively -1 and -2 . We are interested in the case where stable particles with -2 electric charge have substantial relic densities and can capture ${}^4\text{He}^{++}$ nuclei to form a neutral atom. There are three possibilities for this scenario. The first one is to have a relic density of $\bar{U}\bar{U}$, which has -2 charge. For this to be true we should assume that UU is lighter than UD and DD and no processes (apart from electroweak sphalerons) violate the technibaryon number. The second one is to have abundance of ζ that again has -2 charge and the third case is to have both $\bar{U}\bar{U}$ (or DD or $\bar{D}\bar{D}$) and ζ . For the first case to be realized, UU although charged, should be lighter than both UD and DD . This can happen if one assumes that there is an isospin splitting between U and D . This is not hard to imagine since for the same reason in QCD the charged proton is lighter than the neutral neutron. Upon making this assumption, UD and DD will decay through weak interactions to the lightest UU . The technibaryon number is conserved and therefore UU (or $\bar{U}\bar{U}$) is stable. Similarly in

the second case where ζ is the abundant -2 charge particle, ζ must be lighter than ν' and there should be no mixing between the fourth family of leptons and the other three of the Standard Model. The L' number is violated only by sphalerons and therefore after the temperature falls roughly below the electroweak scale Λ_{EW} and the sphalerons freeze out, L' is conserved, which means that the lightest particle, that is ζ in this case, is absolutely stable. We assume also that technibaryons decay to Standard Model particles through Extended Technicolor (ETC) interactions and therefore $TB = 0$. Finally in the third case, we examine the possibility to have both L' and TB conserved after sphalerons have frozen out. In this case, the dark matter would be composed of bound atoms (${}^4He^{++}\zeta^{--}$) and either (${}^4He^{++}(\bar{U}\bar{U})^{--}$) or neutral DD (or $\bar{D}\bar{D}$). We shall examine the three possibilities separately in the next section.

III. THE EXCESS OF THE -2 CHARGED TECHNI-PARTICLES IN THE EARLY UNIVERSE

The calculation of the excess of the technibaryons with respect to the one of the baryons was pioneered in Refs. [31, 32, 33]. In this paper we calculate the excess of $\bar{U}\bar{U}$ and ζ along the lines of [20]. The technicolor and the Standard Model particles are in thermal equilibrium as long as the rate of the weak (and color) interactions is larger than the expansion of the Universe. In addition, the sphalerons allow the violation of the technibaryon number TB , B , L , and L' as long as the temperature of the Universe is higher than roughly Λ_{EW} . It is possible through the equations of thermal equilibrium, sphalerons and overall electric neutrality for the particles of the Universe, to associate the chemical potentials of the various particles. Following [20], we can write down the B , TB , L and L' as

$$B = 12\mu_{uL} + 6\mu_W \quad (1)$$

$$TB = \frac{2}{3}((\sigma_{UU} + \sigma_{UD} + \sigma_{DD})\mu_{UU} + (\sigma_{UD} + 2\sigma_{DD})\mu_W) \quad (2)$$

$$L = 4\mu + 6\mu_W \quad (3)$$

$$L' = 4\sigma_\zeta\mu_{\nu'_L} + 2\sigma_\zeta\mu_W, \quad (4)$$

where μ_{uL} , μ_W , $\mu_{\nu'}$, μ_{UU} are respectively the chemical potentials of the left handed up quark, W , ν' , and UU . μ is the sum of the chemical potentials of the three left handed neutrinos

and the σ_α parameters denote statistical factors for the species α defined as

$$\sigma_\alpha = \frac{6}{4\pi^2} \int_0^\infty dx x^2 \cosh^{-2} \left(\frac{1}{2} \sqrt{x^2 + \left(\frac{m_\alpha}{T^*}\right)^2} \right) \quad \text{for fermions,} \quad (5)$$

$$\sigma_\alpha = \frac{6}{4\pi^2} \int_0^\infty dx x^2 \sinh^{-2} \left(\frac{1}{2} \sqrt{x^2 + \left(\frac{m_\alpha}{T^*}\right)^2} \right) \quad \text{for bosons,} \quad (6)$$

where m_α is the mass of α and T^* is the freeze out temperature for the sphaleron. In the derivation we have assumed for simplicity that the mass of ν' and ζ are very close, so $\sigma_{\nu'} \approx \sigma_\zeta$, and that the Standard Model particles are massless at $T > \Lambda_{EW}$. The sphaleron processes and the condition of the overall electric neutrality impose two extra conditions on the chemical potentials [20]

$$9\mu_{uL} + \frac{3}{2}\mu_{UU} + \mu + \mu_{\nu'} + 8\mu_W = 0, \quad (7)$$

$$Q = 6\mu_{uL} + (2\sigma_{UU} + \sigma_{UD})\mu_{UU} - 2\mu - 6\sigma_\zeta\mu_{\nu'} + (\sigma_{UD} - 4\sigma_\zeta - 18)\mu_W + (14 + \sigma_\zeta)\mu_0 = 0, \quad (8)$$

where μ_0 is the chemical potential of the Higgs boson. Using Eqs. (1), (2), (3), (4), (7), and (8), we can write the ratio of TB/B as a function of the ratios L/B , L'/B and statistical factors as

$$\frac{TB}{B} = -\sigma_{UU} \left(\frac{L'}{B} \frac{1}{3\sigma_\zeta} + 1 + \frac{L}{3B} \right). \quad (9)$$

For the derivation of the above ratio, we have assumed that the electroweak phase transition is of second order, which means that the sphalerons processes freeze out at a temperature slightly lower than the electroweak phase transition. For this reason we have taken $\mu_0 = 0$, since the chemical potential of the Higgs boson in the broken phase should be zero. The calculation in the case of first order phase transition is slightly different, but the results are very similar to the ones of the second order [20]. Furthermore, we have assumed that the mass differences among UU , UD , and DD are not large and therefore we have made the approximation $\sigma_{DD} \simeq \sigma_{UD} \simeq \sigma_{UU}$. In principle we do not have to make this approximation. The ratio (9) would look a bit more complicated but it would not change our physical conclusions. The mass differences among UU , UD , and DD depend on the isospin splitting of the two techniquarks U and D . However, if the splitting is not large, as in the case between up and down quarks in QCD, the mass differences among UU , UD , and DD are small compared to the electroweak energy scale and consequently our approximation is justified. We should emphasize two points regarding the ratio (9). The minus sign in the right hand side denotes the fact that if the quantity inside the parenthesis is positive,

there is an abundance of anti-technibaryons and not technibaryons. In the first case that we investigate, where UU is lighter than UD and DD , an abundance of $\bar{U}\bar{U}$ will provide the charge -2 particles that capture the positively charged nucleus of helium in order to form the neutral dark matter atom. The second point is that (9) seems to diverge if we take the limit where the mass of the ζ and ν' becomes infinite. In that case $\sigma_\zeta \rightarrow 0$ and the ratio diverges. However, L' as seen in (4) depends linearly on σ_ζ and therefore in the limit where the mass of the ζ is very large $L' \rightarrow 0$, unless $\mu_{\nu'} \rightarrow \infty$, which is something unnatural. As we mentioned already there are three different cases that we investigate separately regarding the production of dark matter. The first case is when $\bar{U}\bar{U}$ is the -2 charged particle that will bind with helium to form neutral atom.

A. The case of $\bar{U}\bar{U}$

In this case, $\bar{U}\bar{U}$ is the source of -2 charge particles. The ratio of dark matter produced by the neutral bound state of $({}^4\text{He}^{++}(\bar{U}\bar{U})^{--})$ over the baryon matter is

$$\frac{\Omega_{DM}}{\Omega_B} = \frac{3}{2} \frac{TB}{B} \frac{m_o}{m_p}, \quad (10)$$

where m_o is the mass of the “dark matter atom”, which is approximately the mass of $\bar{U}\bar{U}$ plus 4 GeV (the mass of helium) and m_p is the mass of the proton. In Fig. (1) we show the ratio Ω_{TB}/Ω_B as a function of the mass m of the $\bar{U}\bar{U}$ for several values of the parameter ξ and T^* (the sphaleron freeze out temperature). The parameter ξ is defined as

$$\xi = \frac{L'}{3B\sigma_\zeta} + 1 + \frac{L}{3B}. \quad (11)$$

We should emphasize here that there are two options regarding the new leptons ν' and ζ . If ζ is the lightest between the two and below the electroweak scale no processes violate L' , then ζ will contribute to the relic density of -2 charge particles that bind with helium. We study this case later in subsection C. If ν' is lighter than ζ , nonzero L' could create problems to our model because we have relic density of charged -1 particles. Therefore, we are forced to assume that $L' = 0$. This is a plausible assumption if one allows mixing between ν' and Standard Model leptons, because in that case ν' will decay to lighter leptons and $L' = 0$. Upon making this assumption, i.e. $L' = 0$, $\xi = 1 + L/(3B)$. In the left panel of Fig. (1) we have chosen $T^* = 150$ GeV and several values of ξ ranging from 0.1 to 3. In principle ξ




FIG. 1: *Left Panel:* The ratio of Ω_{TB}/Ω_B as a function of the mass m (in GeV) of the $\bar{U}\bar{U}$ for several values of ξ . The first from the left thin solid line corresponds to $\xi = 0.1$, the thin dashed line to $\xi = 1$, the thick solid line to $\xi = 4/3$, the thick dashed line to $\xi = 2$ and the last solid line to the right corresponds to $\xi = 3$. The horizontal dashed line gives the proper ratio of dark matter over baryon matter, which is approximately 5. For all the curves we have set $T^* = 150$ GeV. *Right Panel:* The same ratio for fixed $\xi = 4/3$, for three different values of the freeze out temperature for the sphalerons T^* , i.e. $T^* = 150$ GeV (thin solid line), $T^* = 200$ GeV (dashed line), and $T^* = 250$ GeV (thick solid line).

can take any positive real value. However, it is logical to assume that the ratio L/B should be around unity. In fact leptogenesis scenarios support a ratio of $L/B = 1$. However, since currently it is not possible to know from observations what is the relic density of the light neutrinos (or antineutrinos), L/B can be also negative. As it can be seen from the figure, the smaller the value of ξ , the lighter becomes the desired $\bar{U}\bar{U}$ that can give the dark matter density. For $\xi = 0$, the excess of technibaryons becomes zero within our approximation. We have plotted several values of ξ , namely $\xi = 0.1, 1, 4/3, 2$ and 3 . In particular, the value $\xi = 4/3$ corresponds to $L = B$. We should stress at this point that for our model to work, ξ should be positive. A negative value of ξ would mean that there is an excess for UU (and not $\bar{U}\bar{U}$), which is positively charged and being bound with ordinary electrons plays a role of an anomalous helium isotope, severely restricted in experimental searches.

B. The case of ζ

Eq. (9) gives the ratio of TB/B as a function of L/B and L'/B . It can be trivially inverted as

$$\frac{L'}{B} = -\sigma_\zeta \left(\frac{3TB}{\sigma_{UU}B} + 3 + \frac{L}{B} \right). \quad (12)$$

In this subsection we investigate the case where ζ is the source of -2 charge particles that can be captured by 4He . For this to be true, ζ must be lighter than ν' and after sphalerons have frozen out, no other processes should violate L' . The term inside the parenthesis of (12) should be negative in order to have abundance of ζ and not anti- ζ . This probably means that a negative ratio L/B is needed. As we mentioned earlier, for our model to be realized, only abundance of -2 charged or neutral particles is accepted. Abundance of charged particles with different charges would cause a serious problem [5, 6, 7, 8, 10, 13]. We study the case where we have abundance for both technibaryons and ζ in the next subsection. Here, we look at the case where $TB = 0$. This can be realized if below the electroweak scale, ETC processes that violate TB exist. In such a case, the lightest technibaryon will decay to lighter Standard Model particles and as a result $TB = 0$. If $TB = 0$, the dark matter density that ζ can provide is given by

$$\frac{\Omega_{L'}}{\Omega_B} = \frac{L'}{B} \frac{m_o}{m_p}, \quad (13)$$

where m_o in this case is the mass of ζ plus the 4 GeV mass of 4He . Fig. (1) shows also the dark matter that ζ with mass m can provide, if we take instead of $\xi = 1 + L/(3B)$, $\xi = -2 - 2L/(3B)$. For example the curve with $\xi = 4/3$, that in the previous case corresponded to $L = B$, now corresponds to $L/B = -5$. This identification is possible because although $\sigma_\zeta(m/T^*)$ and $\sigma_{UU}(m/T^*)$ are defined through Eqs. (5) and (6) respectively, for $m > 500$ GeV (with $T^* = 150$ GeV), the two parameters are approximately equal.

C. The case of ζ plus $\bar{U}\bar{U}$ or DD or $\bar{D}\bar{D}$

The last case we investigate is the one where below the T^* temperature no processes violate TB and L' . This means that the lightest technibaryon and the lightest fourth family lepton are stable objects. In particular, we assume that ζ is lighter than ν' . As for the technibaryons there are two options. The first one is to have DD as the lightest technibaryon.




FIG. 2: *Left Panel:* The value of L/B in terms of the mass of the ζ and the fraction of DD in the dark matter density x , in order the composition of dark matter to be x DD and $(1-x)$ of $({}^4He^{++}\zeta^{--})$. *Right Panel:* Same as in the left panel, for the case where the abundant technibaryon is either $\bar{D}\bar{D}$ or $\bar{U}\bar{U}$. In this case dark matter is made of $(1-x)$ $({}^4He^{++}\zeta^{--})$ and x $\bar{D}\bar{D}$ or $({}^4He^{++}(\bar{U}\bar{U})^{--})$.

In this case, dark matter will be composed of a mixture of neutral DD or $\bar{D}\bar{D}$ and bound $({}^4He^{++}\zeta^{--})$ atoms. The second option is to have UU as the lightest technibaryon. In this case the only acceptable scenario for dark matter is to have a mixture of bound atoms of $({}^4He^{++}\zeta^{--})$ and $({}^4He^{++}(\bar{U}\bar{U})^{--})$. Any other combination, for example an abundance of UU instead of $\bar{U}\bar{U}$ would create problems in our cosmological model since $+2$ charged UU represents a form of anomalous helium. Eq. (12) relates TB/B to L/B and L'/B . If both the technibaryon and ζ contribute to dark matter, the ratio of dark matter over baryon matter is given by

$$\frac{\Omega_{DM}}{\Omega_B} = \frac{\Omega_{TB}}{\Omega_B} + \frac{\Omega_{L'}}{\Omega_B} = 5.09. \quad (14)$$

The contribution of technibaryon is

$$\frac{\Omega_{TB}}{\Omega_B} = \frac{3}{2} \frac{|TB|}{B} \frac{m_{TB}}{m_p}, \quad (15)$$

where m_{TB} is the mass of the lightest technibaryon (plus 4 GeV). We have taken the absolute value of TB because if we have abundance of DD , TB is positive, but for $\bar{U}\bar{U}$ or $\bar{D}\bar{D}$, TB is negative. Similarly by using (12), the contribution of ζ is

$$\frac{\Omega_{L'}}{\Omega_B} = \sigma_\zeta \left| \left(3 + \frac{L}{B} + \frac{3TB}{\sigma_{TB}B} \right) \right| \frac{m_\zeta}{m_p}, \quad (16)$$

where m_ζ is the mass of the ζ (plus 4 GeV). The term inside the absolute brackets should be negative, because we want to have abundance of ζ and not anti- ζ . Therefore we have to

impose the condition

$$3 + \frac{L}{B} + \frac{3TB}{\sigma_{TB}B} < 0. \quad (17)$$

Using the above equations we can rewrite Eq. (14) in a more convenient form

$$\frac{\Omega_{DM}}{\Omega_B} = \left| \sigma_\zeta \left(3 + \frac{L}{B} \right) \frac{m_\zeta}{m_p} \pm \frac{2\sigma_\zeta}{\sigma_{TB}} \frac{m_\zeta}{m_{TB}} 5.09x \right| + 5.09x = 5.09, \quad (18)$$

where x denotes the fraction of dark matter given by the techibaryon. Again, the term inside the absolute brackets should be negative. If the excessive technibaryon is the DD , the above equation should be taken by choosing the plus sign for the term that has \pm . If $\bar{U}\bar{U}$ or $\bar{D}\bar{D}$ is the excessive particle, then the term should be taken with the minus sign.

We first investigate the case of DD mixing. If DD accounts for a component of $x100\%$ of the dark matter density, we can express the ratio L/B as a function of x and the masses of ζ and DD . In particular,

$$\frac{L}{B} = -\frac{5.09m_p}{m_\zeta\sigma_\zeta}[(2z-1)x+1]-3, \quad (19)$$

where $z = (\sigma_\zeta m_\zeta)/(\sigma_{TB}m_{TB})$. If m_ζ is close to m_{TB} , then $z = 1$ (provided that the masses are larger than 350 GeV, if $T^* = 150$ GeV). The study of DD as a dark matter candidate in [20] revealed that DD cannot account for 100% of dark matter density. If we accept that the local dark matter density in the vicinity of the earth is between $0.2-0.4$ GeV/cm³, then with the current exposure of the detectors in CDMS, DD has been ruled out if composes 100% of the dark matter due to its large cross section. However, depending on the mass of DD and the local dark matter density, DD can be a component of dark matter up to 20 – 30% without being excluded by CDMS. This means that in Eq. (19), x should be $0 < x < 0.3$. In Fig. (2), we plot the value of L/B as a function of m_ζ (for $z = 1$) and x (the fraction of dark matter provided by DD), in order the dark matter to be a mixture of x DD and $1-x$ of (${}^4He^{++}\zeta^{--}$). There are three points we would like to stress here. First, it is obvious that the limit $x = 0$ corresponds to the case we studied in the previous subsection. Second, as it can be seen from Fig. (2), L/B gets a large negative value very fast as m_ζ increases beyond 1 – 1.5 TeV. This is something probably unnatural since it is expected that L/B should be of order unity. Third, we see that as x increases, L/B becomes more negative for constant m_ζ . This is because TB is positive and therefore L/B is forced to be more negative in order to get a relic density of ζ . We should also emphasize that if DD is as

low as 200 GeV, then the CDMS constraint becomes tougher. Taking the strict constraint, for low mass for DD , x should be less than 0.1 or even 0.05.

We turn now to the case where TB is negative. This means that the excessive technibaryon is either $\bar{D}\bar{D}$ or $\bar{U}\bar{U}$ according to which one is lighter. We can express as before L/B as

$$\frac{L}{B} = \frac{5.09m_p}{m_\zeta\sigma_\zeta}[(2z+1)x-1]-3. \quad (20)$$

In the right panel of Fig. (2) we show again the projected value of L/B in terms of the mass m_ζ and the fraction x of $\bar{D}\bar{D}$ or $\bar{U}\bar{U}$. We plot x from 0 to 1. The CDMS constraints apply for the $\bar{D}\bar{D}$, but not for $\bar{U}\bar{U}$. This means that if the technibaryon is $\bar{D}\bar{D}$, x should be at most 0.3. From the figure we see that in order L/B to be of order unity, a mass for ζ and $\bar{U}\bar{U}$ between 1 – 2 TeV is needed. Again, the limits $x = 0$ and $x = 1$ were studied in the previous subsections and correspond to the cases of having purely $({}^4He^{++}\zeta^{--})$ or purely $({}^4He^{++}(\bar{U}\bar{U}))^{--}$ respectively.

The case where both UU and ζ are stable techniparticles, offers another possible dark matter scenario. Positively charged UU^{++} and negatively charged ζ^{--} can form “atoms” $(\zeta^{--}UU^{++})$, which behave as cold dark matter species. It resembles the $A^{--}C^{++}$ dark matter atoms of the AC-model [13, 14]. However, as it took place in the AC-cosmology, the existence of +2 charge species, which remain free after all the stages of their binding with –2 charge species and 4He , has a potential danger of anomalous He overproduction. The solution found in the AC-model [13, 14] for this problem of anomalous He involves an additional strict $U(1)$ gauge symmetry, acting on the AC leptons, and a new Coulomb-like long range interaction between the AC-leptons, mediated by the corresponding massless $U(1)$ gauge boson. It could be hardly applied to the case of walking technicolor models. One can have another way to solve this problem. The abundance of free +2 charge techniparticles is suppressed, if all of them are bound with –2 charge techniparticles. Such a complete binding takes place naturally, if the excess of –2 charge techniparticles is larger, than the one of +2 charge techniparticles. Under these conditions, virtually all +2 charge techniparticles are bound in atoms with –2 charge techniparticles, while the residual –2 charge techniparticles bind with 4He in techni-O-helium. The realization of this scenario in the framework of a walking technicolor model and the nontrivial cosmological scenarios involving both cold $(\zeta^{--}UU^{++})$ and warm $({}^4He\zeta^{--})$ forms of composite dark matter goes beyond the scope of the present paper and it is the subject of a separate work.

Due to strong technicolor interactions, the TB excess of $\bar{U}\bar{U}$ suppresses strongly the primordial abundance of the positively charged UU [34]. However in the case of weakly interacting ζ (in the full analogy with the cases of tera-leptons [10] or AC-leptons [13]), its excess does not guarantee a suppression of the corresponding positively charged antiparticles $\bar{\zeta}$. The stages of cosmological evolution resulting in virtually complete elimination of the primordial $\bar{\zeta}$ are stipulated in Appendix 2.

IV. THE CAPTURE OF THE CHARGED TECHNIPARTICLES BY ${}^4\text{He}$

In the Big Bang nucleosynthesis, ${}^4\text{He}$ is formed with an abundance $r_{\text{He}} = 0.1r_B = 8 \cdot 10^{-12}$ and, being in excess, binds all the negatively charged techni-species into atom-like systems. Since the electric charge of $\bar{U}\bar{U}$ (and ζ) is -2 , neutral “atoms” are formed, and $({}^4\text{He}^{++}\zeta^{--})$ “atoms” catalyze effectively the $(\zeta\bar{\zeta})$ binding and annihilation. It turns out [13], that the electromagnetic cascades from this annihilation cannot influence the light element abundance and the energy release of this annihilation takes place so early that it does not distort the CMB spectrum.

At a temperature $T < I_o = Z_{TC}^2 Z_{He}^2 \alpha^2 m_{He}/2 \approx 1.6 \text{ MeV}$, where α is the fine structure constant, and $Z_{TC} = -2$ stands for the electric charge of $\bar{U}\bar{U}$ and/or of ζ , the reaction

$$\zeta^{--} + {}^4\text{He}^{++} \rightarrow \gamma + ({}^4\text{He}\zeta) \quad (21)$$

and/or

$$(\bar{U}\bar{U})^{--} + {}^4\text{He}^{++} \rightarrow \gamma + ({}^4\text{He}(\bar{U}\bar{U})) \quad (22)$$

can take place. In these reactions neutral techni-O-helium “atoms” are produced. The size of these “atoms” is [5, 13]

$$R_o \sim 1/(Z_{TC}Z_{He}\alpha m_{He}) \approx 2 \cdot 10^{-13} \text{ cm} \quad (23)$$

and it can play a nontrivial catalyzing role in nuclear transformations. This aspect needs special thorough study, but some arguments, which we present below following [13], suggest that there should not be contradicting influence on the primordial element abundance.

For our problem another aspect is also important. The reactions (21), and (22) can start only after ${}^4\text{He}$ is formed, which happens at $T < 100 \text{ keV}$. Then, inverse reactions of

ionization by thermal photons support Saha-type relationships between the abundances of these “atoms”, free -2 charged particles, 4He and γ :

$$\frac{n_{He}n_A}{n_\gamma n_{(HeA)}} = \exp\left(-\frac{I_o}{T}\right). \quad (24)$$

From now on, by A^{--} we shall denote $\bar{U}\bar{U}$ and ζ , if the result is independent of their technibaryon or technilepton nature. When T falls below $T_{rHe} \sim I_o/\ln(n_\gamma/n_{He}) \approx I_o/27 \approx 60$ keV, free A^{--} become bound with helium in the reactions of Eqs. (21), and (22). The fraction of free A^{--} , which forms neutral (${}^4He^{++}A^{--}$) depends on the ratio of the abundance of A^{--} over the one of 4He . For $m_\zeta > 50$ GeV and $m_{TB} > 50$ GeV, this ratio is less than 1. Therefore due to 4He excess, all the A^{--} form (${}^4He^{++}A^{--}$) “atoms” through the reactions of Eqs. (21), and (22). Because of this, no free A^{--} are left at the time when $T \sim$ few keV, where $(p^+A^{--})^-$ “ions” or $(p^+p^+A^{--})$ “atoms” could form.

As soon as techni-O-helium (${}^4He^{++}\zeta^{--}$) is formed, heavy antiparticles $\bar{\zeta}$ can penetrate it, expelling 4He and forming ζ -positronium states ($\zeta\bar{\zeta}$), in which the antiparticles annihilate. Therefore the antiparticle $\bar{\zeta}$ can annihilate through formation of positronium, such as

$$(He\zeta) + \bar{\zeta} \rightarrow (\zeta\bar{\zeta} \text{ annihilation products}) + {}^4He. \quad (25)$$

A. 4He capture of free negative charges

At a temperature $T \leq T_{rHe}$, when the reactions of (21), and (22) dominate, the decrease of the free A^{--} abundance due to formation of (${}^4He^{++}A^{--}$) is governed by the equation [13]

$$\frac{dr_A}{dx} = f_{1He} \langle \sigma v \rangle r_A r_{He}, \quad (26)$$

where $x = T/I_o$, $r_{He} = 8 \cdot 10^{-12}$, $\langle \sigma v \rangle$ is given by

$$\langle \sigma v \rangle = \left(\frac{4\pi}{3^{3/2}}\right) \cdot \frac{\bar{\alpha}^2}{I_o \cdot m_{He}} \frac{1}{x^{1/2}},$$

where $\bar{\alpha} = Z_{TC}Z_{He}\alpha$, and (see appendix)

$$f_{1He} \approx m_{Pl}I_o.$$

The solution of Eq. (26) is given by [13]

$$r_A = r_{A0} \exp(-r_{He}J_{He}) = r_{A0} \exp(-1.28 \cdot 10^4),$$

where

$$J_{He} = \int_0^{x_{fHe}} f_{1He} \langle \sigma v \rangle dx =$$

$$= m_{Pl} \left(\frac{4\pi}{3^{3/2}} \right) \cdot \frac{Z_{TC}^2 Z_{He}^2 \alpha^2}{m_{He}} \cdot 2 \cdot \sqrt{x_{fHe}} \approx 1.6 \cdot 10^{15}. \quad (27)$$

$x_{fHe} = 1/27$ is the value of T/I_o where the equilibrium abundance of free charged techniparticles is frozen out and the photodestruction of their atom like bound states with He does not prevent recombination, and m_{Pl} is the Planck mass. Thus, virtually all the free A^{--} are trapped by helium and their remaining abundance becomes exponentially small.

For particles Q^- with charge -1 , as for tera-electrons in the sinister model [9], 4He trapping results in the formation of a positively charged ion $({}^4He^{++}Q^-)^+$, result in dramatic over-production of anomalous hydrogen [10]. Therefore, only the choice of -2 electric charge for stable techniparticles makes it possible to avoid this problem. In this case, 4He trapping leads to the formation of neutral *OLe-helium* “atoms” $({}^4He^{++}A^{--})$, which can catalyze the complete elimination of primordial positively charged anti-technileptons.

B. Complete elimination of antiparticles by techni-O-helium catalysis

For large m_ζ^1 , the primordial abundance of antiparticles $\bar{\zeta}$ is not suppressed. The presence of techni-O-helium $({}^4He^{++}\zeta^{--})$ in this case accelerates the annihilation of these antiparticles through the formation of ζ -positronium. Similar to the case of tera-particles considered in [10] and AC-leptons considered in [13], it can be shown that the products of annihilation cannot cause a back-reaction, ionizing techni-O-helium and suppressing the catalysis.

Indeed, energetic particles, created in $(\zeta\bar{\zeta})$ annihilation, interact with the cosmological plasma. In the development of the electromagnetic cascade, the creation of electron-positron pairs in the reaction $\gamma + \gamma \rightarrow e^+ + e^-$ plays an important role in astrophysical conditions (see [35, 36, 45] for a review). The threshold of this reaction puts an upper limit on the energy of the nonequilibrium photon spectrum in the cascade

$$E_{max} = a \frac{m_e^2}{25T}, \quad (28)$$

where the factor $a = \ln(15\Omega_B + 1) \approx 0.5$ [13].

¹ From now on m_ζ represents the mass of ζ and not the mass of ζ plus 4 GeV.

At a temperature $T > T_{rbHe} = am_e^2/(25I_o) \approx 1 \text{ keV}$, in the spectrum of the electromagnetic cascade from $(\zeta\bar{\zeta})$ annihilation, the maximal energy $E_{max} < I_o$ and the annihilation products cannot ionize (${}^4He^{++}A^{--}$). So, there is no back reaction of the $(\zeta\bar{\zeta})$ annihilation until $T \sim T_{rbHe}$. At that time, practically all free ζ and $\bar{U}\bar{U}$ are bound into (${}^4He^{++}A^{--}$) atoms. For the same reason, electromagnetic showers induced by annihilation products, having a maximal energy below the binding energies of the SBBN nuclei, cannot initiate reactions of non-equilibrium nucleosynthesis and influence the abundance of light elements.

In the absence of back-reaction of annihilation products, nothing prevents the complete elimination of antiparticles $\bar{\zeta}^{++}$ by techni-O-helium catalysis. The $\bar{\zeta}^{++}$ with primordial abundance $r_{\bar{\zeta}}$, can be captured by techni-O-helium (${}^4He^{++}\zeta^{--}$) with abundance $r_{\zeta He} = r_{\zeta} = r_{\bar{\zeta}} + \kappa$ (here the technilepton excess $\kappa = 4 \cdot 10^{-12} f_{\zeta}/S_2$ is given by Eq. (61) of Appendix 2 with $0 \leq f_{\zeta} \leq 1$, being the relative contribution of technileptons into the total dark matter density, and S_2 is the mass of ζ in units of 100 GeV). By definition, $f_{\zeta} = 1 - x$, where x was defined in Eq. (18). The $\bar{\zeta}^{++}$ expels the 4He from the (${}^4He^{++}\zeta^{--}$) and annihilates in ζ -positronium ($\bar{\zeta}^{++}\zeta^{--}$).

The process of $\bar{\zeta}^{++}$ capture by the (${}^4He^{++}\zeta^{--}$) atom looks as follows [13]. Being in thermal equilibrium with the plasma, the free $\bar{\zeta}^{++}$ have momentum $k = \sqrt{2Tm_{\zeta}}$. If their wavelength is much smaller than the size of the (${}^4He^{++}\zeta^{--}$) atom, they can penetrate inside the atom and bind with A^{--} , expelling the 4He from it. The rate of this process is determined by the size of the (${}^4He^{++}\zeta^{--}$) atoms and is given by

$$\langle\sigma v\rangle_0 \sim \pi R_o^2 \sim \frac{\pi}{(\bar{\alpha}m_{He})^2} = \frac{\pi}{2I_o m_{He}} \approx 3 \cdot 10^{-15} \frac{\text{cm}^3}{\text{s}}. \quad (29)$$

Here $\bar{\alpha} = Z_{\zeta}Z_{He}\alpha$. At temperature $T < T_a = \bar{\alpha}^2 m_{He} \frac{m_{He}}{2m_{\zeta}} = \frac{I_o m_{He}}{m_{\zeta}} = 4 \cdot 10^{-2} I_o/S_2$, the wavelength $\lambda = 1/k$ of $\bar{\zeta}^{++}$, exceeds the size $R_o = 1/(\bar{\alpha}m_{\zeta})$ of the (${}^4He^{++}\zeta^{--}$) ‘‘atom’’. The rate of the (${}^4He^{++}\zeta^{--}$) catalysis is suppressed by a factor $(R_o/\lambda)^3 = (T/T_a)^{3/2}$ and is given by

$$\langle\sigma v\rangle_{cat}(T < T_a) = \langle\sigma v\rangle_0 (T/T_a)^{3/2}. \quad (30)$$

The decrease of the antiparticle abundance $r_{\bar{\zeta}}$ is described by

$$\frac{dr_{\bar{\zeta}}}{dx} = f_{1He} \langle\sigma v\rangle r_{\bar{\zeta}}(r_{\bar{\zeta}} + \kappa), \quad (31)$$

where $x = T/I_o$, $r_{He\zeta} = r_{\zeta}$, $\langle\sigma v\rangle$ is given by Eqs. (29) at $T > T_a$ and (30) at $T < T_a$. The

solution of this equation is given in [13] and has the form

$$r_{\bar{\zeta}} = \frac{\kappa \cdot r_{f\bar{\zeta}}}{(\kappa + r_{f\bar{\zeta}}) \exp(\kappa J_o) - r_{f\bar{\zeta}}}, \quad (32)$$

where $r_{f\bar{\zeta}}$ is the frozen concentration of $\bar{\zeta}$ and

$$J_o = \int_0^{x_{fHe}} f_{1He} \langle \sigma v \rangle dx = m_{Pl} \left(\frac{\pi}{2m_{He}} \right) \cdot x_{fHe} \approx 1.4 \cdot 10^{17}, \quad (33)$$

where $x_{fHe} = 1/27$. The factor in the exponent is $\kappa J_o = 6 \cdot 10^5 f_{\zeta}/S_2$. It leads to a huge exponential suppression of the antiparticles at all reasonable values of κ and S_2 .

C. Techni-O-helium in the SBBN

The formation of techni-O-helium reserves a fraction of 4He and thus it changes the primordial abundance of 4He . For the lightest possible masses of the techniparticles $m_{\zeta} \sim m_{TB} \sim 100$ GeV, this effect can reach 50% of the 4He abundance formed in SBBN. Even if the mass of the techniparticles is of the order of TeV, 5% of the 4He abundance is hidden in the techni-O-helium atoms. This can lead to important consequences once we compare the SBBN theoretical predictions to observations.

The question of the participation of techni-O-helium in nuclear transformations and its direct influence on the chemical element production is less evident. Indeed, techni-O-helium looks like an α particle with a shielded electric charge. It can closely approach nuclei due to the absence of a Coulomb barrier. Because of this, it seems that in the presence of techni-O-helium, the character of SBBN processes should change drastically. However, it might not be the case.

The following simple argument can be used to indicate that the techni-O-helium influence on SBBN transformations might not lead to binding of A^{--} with nuclei heavier than 4He . In fact, the size of techni-O-helium is of the order of the size of 4He and for a nucleus A_ZQ with electric charge $Z > 2$, the size of the Bohr orbit for an QA^{--} ion is less than the size of the nucleus A_ZQ . This means that while binding with a heavy nucleus, A^{--} penetrates it and interacts effectively with a part of the nucleus of a size less than the corresponding Bohr orbit. This size corresponds to the size of 4He , making techni-O-helium the most bound QA^{--} atomic state. It favors a picture, according to which a techni-O-helium collision with a nucleus, results in the formation of techni-O-helium and the whole process looks like an elastic collision.

The interaction of the 4He component of $(He^{++}A^{--})$ with a A_ZQ nucleus can lead to a nuclear transformation due to the reaction

$${}^A_ZQ + (HeA) \rightarrow {}^{A+4}_{Z+2}Q + A^{--}, \quad (34)$$

provided that the masses of the initial and final nuclei satisfy the energy condition

$$M(A, Z) + M(4, 2) - I_o > M(A + 4, Z + 2), \quad (35)$$

where $I_o = 1.6$ MeV is the binding energy of techni-O-helium and $M(4, 2)$ is the mass of the 4He nucleus.

This condition is not valid for stable nuclei participating in reactions of the SBBN. However, tritium 3H , which is also formed in SBBN with abundance ${}^3H/H \sim 10^{-7}$ satisfies this condition and can react with techni-O-helium, forming 7Li and opening the path of successive techni-O-helium catalyzed transformations to heavy nuclei. This effect might strongly influence the chemical evolution of matter on the pre-galactic stage and needs a self-consistent consideration within the Big Bang nucleosynthesis network. However, the following arguments show that this effect may not lead to immediate contradiction with observations as it might be expected.

- On the path of reactions (34), the final nucleus can be formed in the excited $(\alpha, M(A, Z))$ state, which can rapidly experience an α -decay, giving rise to techni-O-helium regeneration and to an effective quasi-elastic process of $({}^4He^{++}A^{--})$ -nucleus scattering. It leads to a possible suppression of the techni-O-helium catalysis of nuclear transformations.
- The path of reactions (34) does not stop on 7Li but goes further through ${}^{11}B$, ${}^{15}N$, ${}^{19}F$, ... along the table of the chemical elements.
- The cross section of reactions (34) grows with the mass of the nucleus, making the formation of the heavier elements more probable and moving the main output away from a potentially dangerous Li and B overproduction.

The first publications on possible realistic composite dark matter scenarios [5, 13] gave rise to the development of another aspect of the problem, the Charged massive particles BBN (CBBN), studying the influence of unstable negatively charged massive particles in

BBN [37, 38, 39, 40, 41, 42]. The important difference of CBBN considered in these papers, from our approach, is that singly charged particles X^- with charge -1 do not screen the $+2$ charge of He in a $(HeX)^+$ ion-like bound system, and the Coulomb barrier of the $(HeX)^+$ ion can strongly hamper the path for the creation of isotopes, heavier than 6Li . Therefore, 6Li created in the $D + (HeX)$ reaction cannot dominantly transform into heavier elements and if not destructed by X -decay products, it should remain in the primordial chemical content. It makes the 6Li overproduction found in [37] a really serious trouble for a wide range of parameters for unstable X particles.

It should be noted that the approach of [37] is not supported by [39]. Moreover, we can mention the following effects [13], missed in its solution for the 7Li problem: (i) the competitive process of 7Li creation by a similar mechanism in the reaction ${}^3H + (HeX)^+$ with tritium and (ii) the effects of non-equilibrium nucleosynthesis reactions, induced by hadronic and electromagnetic cascades from products of X decays. The latter effect, which was discussed in [39], implies a self-consistent treatment based on the theory of non-equilibrium cosmological nucleosynthesis [43, 44, 45] (see also [46, 47, 48, 49]). Both effects (i) and (ii) were not studied in [37].

The amount of techni-O-helium in our scenario formally exceeds by a few orders of magnitude the constraint $n_X/s \leq 10^{-17}$, derived for concentration n_X of metastable X^- particles in the units of entropy density s in Eq. (10) of [37]. However, it should be noted that this constraint is not valid for our case if the binding energy $I_o = 1589$ keV of techni-O-helium is taken into account. According to [39], this approximation is valid for $0 < ZZ_{TC}\alpha M_Z R_Z < 1$, where $R_Z \sim 1.2A^{1/3}/200$ MeV $^{-1}$ is the size of nucleus, which is the case for the (HeA) atom. Then the $D + (HeA) \rightarrow {}^6Li + A$ reaction, which the constraint is based on, does not occur. This reaction can take place only if the account for charge distribution in the He nucleus [37] reduces the binding energy of (HeA) down to $E = 1200$ keV or $E = 1150$ keV as discussed in [13]. Then this channel becomes possible, but similar to the case of tritium, the chain of techni-O-helium transformations (34), started from deuterium does not stop on 6Li , but goes further through ${}^{10}B$, ${}^{14}N$, ${}^{18}F$, ... along the table of the chemical elements. Such a qualitative change of the physical picture appeals to necessity in a detailed nuclear physics treatment of the $(A^{--} + \text{nucleus})$ systems and of the whole set of transformations induced by techni-O-helium, including an analysis of possible fast conversion of helium to carbon and of the formation of a $({}^8BeA^{--})$ system, discussed in [13] as potential dangers for our

approach. Though the above arguments do not seem to make these dangers immediate and obvious, a detailed study of this complicated problem is needed.

V. TECHNI-O-HELIUM UNIVERSE

A. Gravitational instability of the techni-O-helium gas

Due to nuclear interactions of its helium constituent with nuclei in cosmic plasma, the techni-O-helium gas is in thermal equilibrium with plasma and radiation on the Radiation Dominance (RD) stage, and the energy and momentum transfer from the plasma is effective. The radiation pressure acting on plasma is then effectively transferred to density fluctuations of techni-O-helium gas and transforms them in acoustic waves at scales up to the size of the horizon. However, as it was first noticed in [5], this transfer to heavy nuclear-interacting species becomes ineffective before the end of the RD stage and such species decouple from plasma and radiation. Consequently, nothing prevents the development of gravitational instability in the gas of these species. This argument is completely applicable to the case of techni-O-helium.

At temperature $T < T_{od} \approx 45S_2^{2/3}$ eV, first estimated in [5] for the case of OLe-helium, the energy and momentum transfer from baryons to techni-O-helium is not effective because $n_B \langle \sigma v \rangle (m_p/m_o)t < 1$, where m_o is the mass of the $tOHe$ atom and $S_2 = \frac{m_o}{100 \text{ GeV}}$. Here

$$\sigma \approx \sigma_o \sim \pi R_o^2 \approx 10^{-25} \text{ cm}^2, \quad (36)$$

and $v = \sqrt{2T/m_p}$ is the baryon thermal velocity. The techni-O-helium gas decouples from the plasma and plays the role of dark matter, which starts to dominate in the Universe at $T_{RM} = 1 \text{ eV}$.

The development of gravitational instabilities of the techni-O-helium gas triggers large scale structure formation, and the composite nature of techni-O-helium makes it more close to warm dark matter.

The total mass of the $tOHe$ gas with density $\rho_d = \frac{T_{RM}}{T_{od}} \rho_{tot}$ within the cosmological horizon $l_h = t$ is

$$M = \frac{4\pi}{3} \rho_d t^3.$$

In the period of decoupling $T = T_{od}$, this mass depends strongly on the techniparticle mass

S_2 and is given by

$$M_{od} = \frac{T_{RM}}{T_{od}} m_{Pl} \left(\frac{m_{Pl}}{T_{od}} \right)^2 \approx 2 \cdot 10^{46} S_2^{-8/3} \text{ g} = 10^{13} S_2^{-8/3} M_{\odot}, \quad (37)$$

where M_{\odot} is the solar mass. The techni-O-helium is formed only at T_{rHe} and its total mass within the cosmological horizon in the period of its creation is $M_o = M_{od}(T_o/T_{od})^3 = 10^{37} \text{ g}$.

On the RD stage before decoupling, the Jeans length λ_J of the $tOHe$ gas was of the order of the cosmological horizon $\lambda_J \sim l_h \sim t$. After decoupling at $T = T_{od}$, it falls down to $\lambda_J \sim v_o t$, where $v_o = \sqrt{2T_{od}/m_o}$. Though after decoupling the Jeans mass in the $tOHe$ gas correspondingly falls down

$$M_J \sim v_o^3 M_{od} \sim 3 \cdot 10^{-14} M_{od},$$

one should expect strong suppression of fluctuations on scales $M < M_o$, as well as adiabatic damping of sound waves in the RD plasma for scales $M_o < M < M_{od}$. It provides suppression of small scale structure in the considered model for all reasonable masses of techniparticles.

The cross section of mutual collisions of techni-O-helium “atoms” is given by Eq. (36). The $tOHe$ “atoms” can be considered as collision-less gas in clouds with a number density n_o and a size R , if $n_o R < 1/\sigma_o$. This condition is valid for the techni-O-helium gas in galaxies.

Mutual collisions of techni-O-helium “atoms” determine the evolution timescale for a gravitationally bound system of collision-less $tOHe$ gas

$$t_{ev} = 1/(n\sigma_o v) \approx 2 \cdot 10^{20} (1 \text{ cm}^{-3}/n)^{7/6} \text{ s},$$

where the relative velocity $v = \sqrt{GM/R}$ is taken for a cloud of mass M_o and an internal number density n . This timescale exceeds substantially the age of the Universe and the internal evolution of techni-O-helium clouds cannot lead to the formation of dense objects. Being decoupled from baryonic matter, the $tOHe$ gas does not follow the formation of baryonic astrophysical objects (stars, planets, molecular clouds...) and forms dark matter halos of galaxies.

B. Techniparticle component of cosmic rays

The nuclear interaction of techni-O-helium with cosmic rays gives rise to ionization of this bound state in the interstellar gas and to acceleration of free techniparticles in the Galaxy.

During the lifetime of the Galaxy $t_G \approx 3 \cdot 10^{17}$ s, the integral flux of cosmic rays

$$F(E > E_0) \approx 1 \cdot \left(\frac{E_0}{1 \text{ GeV}} \right)^{-1.7} \text{ cm}^{-2} \text{ s}^{-1}$$

can disrupt the fraction of galactic techni-O-helium $\sim F(E > E_{min})\sigma_o t_G \leq 10^{-3}$, where we took $E_{min} \sim I_o$. Assuming a universal mechanism of cosmic ray acceleration, a universal form of their spectrum, taking into account that the ${}^4\text{He}$ component corresponds to $\sim 5\%$ of the proton spectrum, and that the spectrum is usually reduced to the energy per nucleon, the anomalous low $Z/A - 2$ charged techniparticle component can be present in cosmic rays at a level of

$$\frac{A^{--}}{He} \geq 3 \cdot 10^{-7} \cdot S_2^{-3.7}. \quad (38)$$

This flux may be within the reach for PAMELA and AMS02 cosmic ray experiments.

Recombination of free techniparticles with protons and nuclei in the interstellar space can give rise to radiation in the range from few tens of keV - 1 MeV. However such a radiation is below the cosmic nonthermal electromagnetic background radiation observed in this range.

C. Effects of techni-O-helium catalyzed processes in the Earth

The first evident consequence of the proposed excess is the inevitable presence of $tOHe$ in terrestrial matter. This is because terrestrial matter appears opaque to $tOHe$ and stores all its in-falling flux.

If the $tOHe$ capture by nuclei is not effective, its diffusion in matter is determined by elastic collisions, which have a transport cross section per nucleon

$$\sigma_{tr} = \pi R_o^2 \frac{m_p}{m_o} \approx 10^{-27} / S_2 \text{ cm}^2. \quad (39)$$

In atmosphere, with effective height $L_{atm} = 10^6$ cm and baryon number density $n_B = 6 \cdot 10^{20} \text{ cm}^{-3}$, the opacity condition $n_B \sigma_{tr} L_{atm} = 6 \cdot 10^{-1} / S_2$ is not strong enough. Therefore, the in-falling $tOHe$ particles are effectively slowed down only after they fall down terrestrial surface in $16S_2$ meters of water (or $4S_2$ meters of rock). Then they drift with velocity $V = \frac{g}{n\sigma v} \approx 8S_2 A^{1/2} \text{ cm/s}$ (where $A \sim 30$ is the average atomic weight in terrestrial surface matter, and $g = 980 \text{ cm/s}^2$), sinking down the center of the Earth on a timescale $t = R_E / V \approx 1.5 \cdot 10^7 S_2^{-1}$ s, where R_E is the radius of the Earth.

The in-falling techni-O-helium flux from dark matter halo is $\mathcal{F} = n_o v_h / 8\pi$, where the number density of $tOHe$ in the vicinity of the Solar System is $n_o = 3 \cdot 10^{-3} S_2^{-1} \text{ cm}^{-3}$ and the averaged velocity $v_h \approx 3 \cdot 10^7 \text{ cm/s}$. During the lifetime of the Earth ($t_E \approx 10^{17} \text{ s}$), about $2 \cdot 10^{38} S_2^{-1}$ techni-O-helium atoms were captured. If $tOHe$ dominantly sinks down the Earth, it should be concentrated near the Earth's center within a radius $R_{oc} \sim \sqrt{3T_c / (m_o 4\pi G \rho_c)}$, which is $\leq 10^8 S_2^{-1/2} \text{ cm}$, for the Earth's central temperature $T_c \sim 10^4 \text{ K}$ and density $\rho_c \sim 4 \text{ g/cm}^3$.

Near the Earth's surface, the techni-O-helium abundance is determined by the equilibrium between the in-falling and down-drifting fluxes. It gives

$$n_{oE} = 2\pi\mathcal{F}/V = 3 \cdot 10^3 \cdot S_2^{-2} \cdot A^{-1/2} \text{ cm}^{-3},$$

or for $A \sim 30$ about $5 \cdot 10^2 \cdot S_2^{-2} \text{ cm}^{-3}$. This number density corresponds to the fraction

$$f_{oE} \sim 5 \cdot 10^{-21} \cdot S_2^{-2}$$

relative to the number density of the terrestrial atoms $n_A \approx 10^{23} \text{ cm}^{-3}$.

These neutral (${}^4He^{++}A^{--}$) “atoms” may provide a catalysis of cold nuclear reactions in ordinary matter (much more effectively than muon catalysis). This effect needs a special and thorough investigation. On the other hand, if A^{--} capture by nuclei, heavier than helium, is not effective and does not lead to a copious production of anomalous isotopes, the (${}^4He^{++}A^{--}$) diffusion in matter is determined by the elastic collision cross section (39) and may effectively hide techni-O-helium from observations.

One can give the following argument for an effective regeneration and quasi-elastic collisions of techni-O-helium in terrestrial matter. The techni-O-helium can be destroyed in the reactions (34). Then, free A^{--} are released and due to a hybrid Auger effect (capture of A^{--} , ejection of ordinary e from the atom with atomic number A , and charge of the nucleus Z), A^{--} -atoms are formed, in which A^{--} occupies highly an excited level of the (A_ZQA) system, which is still much deeper than the lowest electronic shell of the considered atom. The (A_ZQA) atomic transitions to lower-lying states cause radiation in the intermediate range between atomic and nuclear transitions. In course of this falling down to the center of the ($Z - A^{--}$) system, the nucleus approaches A^{--} . For $A > 3$ the energy of the lowest state n (given by $E_n = \frac{M\bar{\alpha}^2}{2n^2} = \frac{2Am_p Z^2 \alpha^2}{n^2}$) of the ($Z - A^{--}$) system (having reduced mass $M \approx Am_p$) with a Bohr orbit $r_n = \frac{n}{M\bar{\alpha}} = \frac{n}{2AZm_p\alpha}$, exceeding the size of the nucleus $r_A \sim A^{1/3}m_\pi^{-1}$

(m_π being the mass of the pion), is less than the binding energy of $tOHe$. Therefore the regeneration of techni-O-helium in a reaction, inverse to (34), takes place. An additional reason for the domination of the elastic channel of the reactions (34) is that the final state nucleus is created in the excited state and its de-excitation via α -decay can also result in techni-O-helium regeneration. If regeneration is not effective and A^{--} remains bound to the heavy nucleus, anomalous isotope of $Z - 2$ element should appear. This is a serious problem for the considered model.

However, if the general picture of sinking down is valid, it might give no more than the ratio $f_{oE} \sim 5 \cdot 10^{-21} \cdot S_2^{-2}$ of number density of anomalous isotopes to the number density of atoms of terrestrial matter around us, which is below the experimental upper limits for elements with $Z \geq 2$. For comparison, the best upper limits on the anomalous helium were obtained in [50]. It was found, by searching with the use of laser spectroscopy for a heavy helium isotope in the Earth's atmosphere, that in the mass range 5 GeV - 10000 GeV, the terrestrial abundance (the ratio of anomalous helium number to the total number of atoms in the Earth) of anomalous helium is less than $2 \cdot 10^{-19} - 3 \cdot 10^{-19}$.

D. Direct search for techni-O-helium

It should be noted that the nuclear cross section of the techni-O-helium interaction with matter escapes the severe constraints [24] on strongly interacting dark matter particles (SIMPs) [23, 24] imposed by the XQC experiment [51].

In underground detectors, $tOHe$ “atoms” are slowed down to thermal energies and give rise to energy transfer $\sim 2.5 \cdot 10^{-3} \text{ eV} A / S_2$, far below the threshold for direct dark matter detection. It makes this form of dark matter insensitive to the CDMS constraints. However, $tOHe$ induced nuclear transformation can result in observable effects.

Therefore, a special strategy of such a search is needed, that can exploit sensitive dark matter detectors on the ground or in space. In particular, as it was revealed in [53], a few g of superfluid 3He detector [52], situated in ground-based laboratory can be used to put constraints on the in-falling techni-O-helium flux from the galactic halo.

VI. DISCUSSION

In this paper we explored the cosmological implications of a walking technicolor model with doubly charged technibaryons and technileptons. The considered model escapes most of the drastic problems of the Sinister Universe [9], related to the primordial 4He cage for -1 charge particles and a consequent overproduction of anomalous hydrogen [10]. These charged 4He cages pose a serious problem for composite dark matter models with single charged particles, since their Coulomb barrier prevents successful recombination of positively and negatively charged particles. The doubly charged A^{--} techniparticles considered in this paper, bind with 4He in the techni-O-helium neutral states.

To avoid overproduction of anomalous isotopes, an excess of -2 charged techniparticles over their antiparticles should be generated in the Universe. In all the previous realizations of composite dark matter scenarios, this excess was put by hand to saturate the observed dark matter density. In our paradigm, this abundance of techibaryons and/or technileptons is connected naturally to the baryon relic density.

A challenging problem that we leave for future work is the nuclear transformations, catalyzed by techni-O-helium. The question about their consistency with observations remains open, since special nuclear physics analysis is needed to reveal what are the actual techni-O-helium effects in SBBN and in terrestrial matter. Another aspect of the considered approach is more clear. For reasonable values of the techiparticle mass, the amount of primordial 4He , bound in this atom like state is significant and should be taken into account in comparison to observations.

The destruction of techni-O-helium by cosmic rays in the Galaxy releases free charged techniparticles, which can be accelerated and contribute to the flux of cosmic rays. In this context, the search for techniparticles at accelerators and in cosmic rays acquires the meaning of a crucial test for the existence of the basic components of the composite dark matter. At accelerators, techniparticles would look like stable doubly charged heavy leptons, while in cosmic rays, they represent a heavy -2 charge component with anomalously low ratio of electric charge to mass.

To conclude, walking technicolor cosmology can naturally resolve most of problems of composite dark matter. Therefore, the model considered in this paper with stable -2 charged particles might provide a realistic physical basis for a composite dark matter scenario.

Acknowledgements

The work of C.K. is supported by the Marie Curie Fellowship under contract MEIF-CT-2006-039211. We are grateful to K.M. Belotsky and F.Sannino for reading the manuscript and for important remarks. M.Kh. is grateful to CERN (Geneve, Switzerland) and to ICTP (Trieste, Italy) for hospitality.

Appendix 1. Charge asymmetry in freezing out of particles and antiparticles

The frozen number density of cosmic relics, which were in equilibrium with the primordial plasma, is conventionally deduced² from the equation [54]

$$\dot{n} + 3Hn = \langle \sigma_{ann} v \rangle (n_{eq}^2 - n^2). \quad (40)$$

This equation is written for the case of a charge symmetry of the particles in question, i.e. for the case when number densities of particles X and antiparticles \bar{X} are equal $n_X = n_{\bar{X}} = n$. The value n_{eq} corresponds to their equilibrium number density and is given by the Boltzmann distribution

$$n_{eq} = g_S \left(\frac{mT}{2\pi} \right)^{3/2} \exp \left(-\frac{m}{T} \right). \quad (41)$$

Here g_S and m are the number of spin states and the mass of the given particle.

During the cooling, n_{eq} decreases exponentially and becomes, below the freezing out temperature T_f , much smaller than the real density n , so the term $\langle \sigma_{ann} v \rangle n_{eq}^2$, describing the creation of $X\bar{X}$ from the plasma can be neglected [55]. It allows to obtain an approximate solution of Eq. (40).

In case of a charge asymmetry one needs to split Eq. (40) in two: for n_X and $n_{\bar{X}}$, which are not equal now.

$$\begin{aligned} \dot{n}_X + 3Hn_X &= \langle \sigma_{ann} v \rangle (n_{eqX} n_{eq\bar{X}} - n_X n_{\bar{X}}), \\ \dot{n}_{\bar{X}} + 3Hn_{\bar{X}} &= \langle \sigma_{ann} v \rangle (n_{eqX} n_{eq\bar{X}} - n_X n_{\bar{X}}). \end{aligned} \quad (42)$$

The values n_{eqX} and $n_{eq\bar{X}}$ are given by Eq. (41) with inclusion of the chemical potential, which for X and for \bar{X} are related as $\mu_X = -\mu_{\bar{X}} = \mu$ (see, e.g., [56]). So

$$n_{eqX, \bar{X}} = \exp \left(\pm \frac{\mu}{T} \right) n_{eq}, \quad (43)$$

² We follow here the results obtained in [10, 13] with the help of K.M. Belotsky

where upper and lower signs are for X and \bar{X} respectively. So

$$n_{eq\,X}n_{eq\,\bar{X}} = n_{eq}^2. \quad (44)$$

A degree of asymmetry will be described in the conventional manner (as for baryons) by the ratio of the difference between n_X and $n_{\bar{X}}$ to the number density of relic photons at the modern period

$$\kappa_{\gamma\,mod} = \frac{n_{X\,mod} - n_{\bar{X}\,mod}}{n_{\gamma\,mod}}. \quad (45)$$

However, for practical purposes it is more suitable to use the ratio to the entropy density, which unlike Eq. (45), does not change in time provided entropy conservation. The photon number density n_γ and the entropy density s are given by

$$n_\gamma = \frac{2\zeta(3)}{\pi^2}T^3, \quad s = \frac{2\pi^2 g_s}{45}T^3 = 1.80g_s n_\gamma, \quad (46)$$

where

$$g_s = \sum_{bos} g_s \left(\frac{T_{bos}}{T}\right)^3 + \frac{7}{8} \sum_{ferm} g_s \left(\frac{T_{ferm}}{T}\right)^3. \quad (47)$$

The sums in Eq. (47) are over ultrarelativistic bosons and fermions. So

$$\kappa = \frac{n_X - n_{\bar{X}}}{s}, \quad \kappa = \frac{\kappa_{\gamma\,mod}}{1.8g_{s\,mod}}, \quad (48)$$

where $g_{s\,mod} = 43/11 \approx 3.91$. Eq. (48) provides a connection between n_X and $n_{\bar{X}}$. Let us pass to the variables

$$r_+ = \frac{n_X}{s}, \quad r_- = \frac{n_{\bar{X}}}{s}, \quad r = \frac{n_X + n_{\bar{X}}}{s}, \quad x = \frac{T}{m}. \quad (49)$$

The apparent relations between the r_i are

$$r_+ - r_- = \kappa, \quad r_+ + r_- = r. \quad (50)$$

Provided that the essential entropy redistribution does not take place ($g_s = \text{const.}$) during the period of freezing out, a transformation to the variable x is possible

$$-Hdt = dT/T = dx/x.$$

On the RD stage the Hubble parameter depends on T as

$$H = \frac{2\pi}{3} \sqrt{\frac{\pi g_\epsilon}{5}} \frac{T^2}{m_{Pl}}, \quad (51)$$

where g_ϵ is given by

$$g_\epsilon = \sum_{bos} g_S \left(\frac{T_{bos}}{T} \right)^4 + \frac{7}{8} \sum_{ferm} g_S \left(\frac{T_{ferm}}{T} \right)^4. \quad (52)$$

For r_+ , r_- and r from Eqs. (42) one obtains the equations

$$\begin{aligned} \frac{dr_+}{dx} &= f_1 \langle \sigma_{ann} v \rangle (r_+ (r_+ - \kappa) - f_2(x)) \\ \frac{dr_-}{dx} &= f_1 \langle \sigma_{ann} v \rangle (r_- (r_- + \kappa) - f_2(x)) \\ \frac{dr}{dx} &= \frac{1}{2} f_1 \langle \sigma_{ann} v \rangle (r^2 - \kappa^2 - 4f_2(x)). \end{aligned} \quad (53)$$

Here

$$\begin{aligned} f_1 &= \frac{s}{Hx}, \\ f_2(x) &= \frac{n_{eq}^2}{s^2} = \frac{45^2 g_S^2}{2^5 \pi^7 g_s^2 x^3} \exp \left(-\frac{2}{x} \right). \end{aligned} \quad (54)$$

By using Eqs. (46) and Eq. (51), one finds that on the RD stage f_1 is

$$f_1 = \sqrt{\frac{\pi g_s^2}{45 g_\epsilon}} m_{Pl} m$$

and independent of x .

To solve Eqs. (53) analogously to Eq. (40), namely neglecting $f_2(x)$ in them, starting with some $x = x_f$, it would not be more difficult to define the moment $x = x_f$. Nonetheless, if one supposes that such a moment is defined, r_i will be

$$\begin{aligned} r_+(x \approx 0) &= \frac{\kappa \cdot r_{+f}}{r_{+f} - (r_{+f} - \kappa) \exp(-\kappa J)}, \\ r_-(x \approx 0) &= \frac{\kappa \cdot r_{-f}}{(\kappa + r_{-f}) \exp(\kappa J) - r_{-f}}, \\ r(x \approx 0) &= \kappa \frac{(\kappa + r_f) \exp(\kappa J) + r_f - \kappa}{(\kappa + r_f) \exp(\kappa J) - (r_f - \kappa)}. \end{aligned} \quad (55)$$

Here $r_{if} = r_i(x = x_f)$,

$$J = \int_0^{x_f} f_1 \langle \sigma_{ann} v \rangle dx.$$

All r_i (at any moment) are related with the help of Eqs. (50). Taking into account Eq. (43) or Eq. (44) for r_{if} one obtains

$$\begin{aligned} r_{\pm f} &= \frac{1}{2} \left(\sqrt{4f_2(x_f) + \kappa^2} \pm \kappa \right), \\ r_f &= \sqrt{4f_2(x_f) + \kappa^2}. \end{aligned} \quad (56)$$

For $\langle\sigma_{ann}v\rangle$ independent of x on the RD stage, f_1 also independent of x , and x_f defined from the condition $R(T_f) = H(T_f)$ for the reaction rate $R(T_f) = n_{eq}(T_f) \langle\sigma_{ann}v(T_f)\rangle$, leading to

$$\begin{aligned} n_{eq}(T_f) \langle\sigma_{ann}v(T_f)\rangle / H(T_f) &= \frac{n_{eq}}{s} \cdot \frac{s}{Hx_f} \cdot \langle\sigma_{ann}v(x_f)\rangle \cdot x_f = \\ &= \sqrt{f_2(x_f)f_1} \langle\sigma_{ann}v(x_f)\rangle \cdot x_f = 1, \end{aligned} \quad (57)$$

one obtains

$$\sqrt{f_2(x_f)} = \frac{1}{f_1 \langle\sigma_{ann}v\rangle \cdot x_f} = \frac{1}{J}. \quad (58)$$

If (a) $\langle\sigma_{ann}v\rangle = \alpha^2/m^2$ or (b) $\langle\sigma_{ann}v\rangle = \alpha/\sqrt{Tm^3}$ and one assumes $f_1 = \text{const}$, then

$$\begin{aligned} J_a &= \sqrt{\frac{\pi g_s^2}{45g_\epsilon} m_{Pl} \frac{\alpha^2}{m}} x_f, \\ J_b &= \sqrt{\frac{\pi g_s^2}{45g_\epsilon} m_{Pl} \frac{\alpha}{m}} 2\sqrt{x_f}. \end{aligned} \quad (59)$$

Appendix 2. Primordial technileptons from the Big Bang Universe

As already mentioned, the minimal walking technicolor model considered in this paper can allow the creation of ζ excess that might contribute (or even saturate) to the modern dark matter density in the form of techni-O-helium “atoms”. For light baryon excess $\eta_B = n_{B\text{mod}}/n_{\gamma\text{mod}} = 6 \cdot 10^{-10}$, it gives a ζ -excess

$$\eta_\zeta = n_{\zeta\text{mod}}/n_{\gamma\text{mod}} = 3 \cdot 10^{-11} f_\zeta \left(\frac{100 \text{ GeV}}{m_\zeta} \right), \quad (60)$$

where m_ζ is the mass of ζ . By definition f_ζ is the contribution of ζ into the modern dark matter density $f_\zeta = 1 - x$, where x was defined in Eq. (18). For future use, following [9, 10], it is convenient to relate the baryon density $\Omega_B = 0.044$ and the technilepton density $\Omega_{L'} = 0.224$ with the entropy density s , and to introduce $r_B = n_B/s$ and $r_\zeta = n_\zeta/s$. Taking into account that $s_{\text{mod}} = 7.04 \cdot n_{\gamma\text{mod}}$, one obtains $r_B \sim 8 \cdot 10^{-11}$ and

$$r_\zeta = 4 \cdot 10^{-12} \left(\frac{100 \text{ GeV}}{m_\zeta} \right) = 4 \cdot 10^{-12} f_\zeta / S_2. \quad (61)$$

Chronological cornerstones of the technilepton-Universe

After the generation of technilepton asymmetry, the thermal history of technileptons in chronological order looks as follows:

1) $10^{-10}S_2^{-2}\text{s} \leq t \leq 6 \cdot 10^{-8}S_2^{-2}\text{s}$ at $m_\zeta \geq T \geq T_f = m_\zeta/31 \approx 3S_2\text{ GeV}$. ζ -lepton pair $\zeta\bar{\zeta}$ annihilation and freezing out. For large m_ζ the abundance of frozen out ζ -lepton pairs is not suppressed in spite of an ζ -lepton excess.

2) $t \sim 2.4 \cdot 10^{-3}S_2^{-2}\text{s}$ at $T \sim I_\zeta = 20S_2\text{ MeV}$. The temperature corresponds to the binding energy $I_\zeta = Z_\zeta^4\alpha^2m_\zeta/4 \approx 20S_2\text{ MeV}$ ($Z_\zeta = 2$) of ζ -positronium “atoms” ($\zeta^{--}\bar{\zeta}^{++}$), in which $\bar{\zeta}^{++}$ annihilate. At large m_ζ this annihilation is not at all effective to reduce the $\zeta\bar{\zeta}$ pairs abundance. These pairs are eliminated in the course of the successive evolution of ζ -matter.

3) $100\text{ s} \leq t \leq 300\text{ s}$ at $100\text{ keV} \geq T \geq I_o/27 \approx 60\text{ keV}$. 4He is formed in the SBBN and virtually all free ζ^{--} are trapped by 4He in (${}^4He^{++}\zeta^{--}$). Note that in the period $100\text{ keV} \leq T \leq 1.6\text{ MeV}$, 4He is not formed, therefore it is only after *the first three minutes*, when (${}^4He^{++}\zeta^{--}$) trapping of ζ^{--} can take place. Being formed, techni-OLe-helium catalyzes the binding of free $\bar{\zeta}^{++}$ with its constituent ζ^{--} into ζ -positronium and complete annihilation of all the primordial antiparticles. At large m_ζ , effects of ($\zeta^{--}\bar{\zeta}^{++}$) annihilation, catalyzed by techni-O-helium, do not cause any contradictions with observations.

4) $t \sim 10^{12}\text{ s}$ at $T \sim T_{RM} \approx 1\text{ eV}$. The techniparticle dominance starts with techni-O-helium “atoms” playing the role more close to warm dark matter in the formation of large scale structures.

Freezing out of ζ -leptons

In the early Universe at temperatures highly above their masses, the ζ -fermions were in thermodynamical equilibrium with the relativistic plasma. It means that at $T > m_\zeta$ the excessive ζ were accompanied by $\zeta\bar{\zeta}$ pairs.

During the expansion, when the temperature T falls below the mass of the ζ -particles, the concentration of particles and antiparticles is given by the equilibrium. The equilibrium concentration of $\zeta\bar{\zeta}$ pairs starts to decrease at $T < m_\zeta = 100S_2\text{ GeV}$. At the freezing out temperature T_f , the rate of expansion exceeds the rate of annihilation to photons $\zeta\bar{\zeta} \rightarrow \gamma\gamma$, to W, Z -bosons $\zeta\bar{\zeta} \rightarrow WW(ZZ)$, or to pairs of light fermions f (quarks and charged leptons) $\zeta\bar{\zeta} \rightarrow \bar{f}f$ (the latter takes place both due to electromagnetic and weak interactions). Then ζ leptons and their antiparticles $\bar{\zeta}$ are frozen out.

In the case of freezing out of ζ -leptons one has (see Appendix 1)

$$f_{1\zeta} = \sqrt{\frac{\pi g_s^2}{45g_\epsilon}} m_{Pl} m_\zeta \approx 2.5 m_{Pl} m_\zeta,$$

$$\langle \sigma_{ann} v \rangle = \frac{\bar{\alpha}^2}{m_\zeta^2} \text{ and}$$

$$J_\zeta = \sqrt{\frac{\pi g_s^2}{45g_\epsilon}} m_{Pl} \frac{\bar{\alpha}^2}{m_\zeta} x_f, \quad (62)$$

where $\bar{\alpha} = Z_\zeta^2 \alpha + \bar{\alpha}_{ew}$, $Z_\zeta = 2$ is the absolute value of the electric charge of the ζ and $\bar{\alpha}_{ew}$ takes into account the effects of W and Z bosons in technilepton annihilation. By putting in Eq. (54) $g_s = 2$, $g_s \sim 100$, one obtains the solution of the transcendent equation (58)

$$\begin{aligned} x_f &\approx \left(\ln \left(\frac{45g_s}{2^{5/2}\pi^{7/2}g_s} \cdot f_{1\zeta} \langle \sigma_{ann} v \rangle \right) \right)^{-1} \approx \\ &\approx \frac{1}{30} \cdot \frac{1}{(1 - \ln(S_2)/30)}. \end{aligned}$$

Taking $g_s \approx g_\epsilon \sim 100$, one finds from Eq. (62) $J_\zeta = 6.5 \cdot 10^{13}/S_2(1 - \ln(S_2)/30)^{-1}$ and from Eq. (58) $\sqrt{4f_2(x_f)} = 2/J_\zeta = 3 \cdot 10^{-13} S_2 \cdot (1 - \ln(S_2)/30)$. For $\kappa = r_\zeta = 4 \cdot 10^{-12} f_\zeta/S_2$, one has $\kappa J_\zeta = 26 f_\zeta/S_2^2$. At $S_2 < 2.7$ $4f_2(x_f) < \kappa^2$ and $r_{\pm f}$ is given by Eq. (56). Since $4f_2(x_f) \gg \kappa^2$ for $S_2 \gg 1$, one obtains from Eq. (56)

$$r_{\pm f} = \frac{1}{2} \left(\sqrt{4f_2(x_f)} \pm \kappa \right). \quad (63)$$

The frozen out abundances of ζ -leptons and their antiparticles are given by

$$\begin{aligned} r_\zeta &= \frac{\kappa \cdot r_{+f}}{r_{+f} - (r_{+f} - \kappa) \exp(-\kappa J_\zeta)} = F_\zeta(S_2), \\ r_{\bar{\zeta}} &= \frac{\kappa \cdot r_{-f}}{(\kappa + r_{-f}) \exp(\kappa J_\zeta) - r_{-f}} = F_{\bar{\zeta}}(S_2). \end{aligned} \quad (64)$$

For growing $S_2 \gg 1$, the solution Eq. (64) approaches the values

$$\begin{aligned} r_\zeta &\approx \sqrt{f_2(x_f)} + \kappa/2 \approx \\ &\approx 1.5 \cdot 10^{-13} S_2 \cdot (1 - \ln(S_2)/30) + 2 \cdot 10^{-12} f_\zeta/S_2, \\ r_{\bar{\zeta}} &\approx \sqrt{f_2(x_f)} - \kappa/2 \approx \\ &\approx 1.5 \cdot 10^{-13} S_2 \cdot (1 - \ln(S_2)/30) - 2 \cdot 10^{-12} f_\zeta/S_2. \end{aligned} \quad (65)$$

At $S_2 < 5f_\zeta$, the factor in the exponent κJ_ζ exceeds 1, and some suppression of the $(\bar{\zeta})$ -abundance takes place. For S_2 close to 1, one has

$$\begin{aligned} r_\zeta &= F_\zeta(S_2) \approx \kappa = 4 \cdot 10^{-12} f_\zeta/S_2, \\ r_{\bar{\zeta}} &= F_{\bar{\zeta}}(S_2) \approx 5 \cdot 10^{-3} \kappa S_2^4 \exp(-26 f_\zeta/S_2^2). \end{aligned} \quad (66)$$

At $S_2 = 1$, the factor in the exponent reaches the value $\kappa J_\zeta = 26f_\zeta$ and the solution Eq. (64) gives $r_\zeta \approx \kappa = 4 \cdot 10^{-12}f_\zeta$ and

$$\begin{aligned} r_{\bar{\zeta}} &\approx \frac{\kappa \cdot r_{-f}}{\kappa + r_{-f}} \exp(-\kappa J_\zeta) \approx r_{-f} \exp(-\kappa J_\zeta) \\ &\approx 10^{-14} \exp(-26f_\zeta) \approx 3 \cdot 10^{-25} \end{aligned}$$

for $f_\zeta = 1$ and $r_{-f} \approx 10^{-14}$ from Eq. (56).

The S_2 -dependence of the frozen out abundances (in units of the entropy density) of the ζ leptons and their antiparticles are

$$\begin{aligned} r_\zeta &= F_\zeta(S_2), \\ r_{\bar{\zeta}} &= F_{\bar{\zeta}}(S_2), \end{aligned} \tag{67}$$

given by Eq. (64). For growing $S_2 \gg 1$, the solution Eq. (64) approaches the values

$$\begin{aligned} r_{AC} &\approx \sqrt{f_2(x_f)} + \kappa/2 \approx \\ &\approx 1.5 \cdot 10^{-13} S_2 \cdot (1 - \ln(S_2)/30) + 2 \cdot 10^{-12} f_\zeta / S_2, \\ r_{\bar{AC}} &\approx \sqrt{f_2(x_f)} - \kappa/2 \approx \\ &\approx 1.5 \cdot 10^{-13} S_2 \cdot (1 - \ln(S_2)/30) - 2 \cdot 10^{-12} f_\zeta / S_2. \end{aligned} \tag{68}$$

At $S_2 < 5f_\zeta$, there is an exponential suppression of the $\bar{\zeta}$ abundance. For S_2 close to 1, one has

$$\begin{aligned} r_\zeta &= F_\zeta(S_2) \approx \kappa = 4 \cdot 10^{-12} / S_2, \\ r_{\bar{\zeta}} &= f_{\bar{\zeta}}(S_2) \approx 5 \cdot 10^{-3} \kappa S_2^4 \exp(-26f_\zeta / S_2^2). \end{aligned} \tag{69}$$

At $S_2 = 1$, the solution Eq. (69) gives

$$r_\zeta \approx \kappa_A = \kappa_C = 2 \cdot 10^{-12} f_\zeta,$$

and

$$r_{\bar{\zeta}} \approx 3 \cdot 10^{-25}.$$

On the other hand at $S_2 > 5$, the concentration of frozen out ζ -lepton pairs exceeds the one of the ζ -lepton excess given by Eq. (61) and this effect grows with S_2 as $\propto S_2^2$ at large S_2 . So in this moment, in spite of an assumed ζ -lepton asymmetry, the frozen out concentration of

antiparticles $\bar{\zeta}$ is not strongly suppressed and they cannot be neglected in the cosmological evolution of ζ -matter.

The antiparticles $\bar{\zeta}$ should be effectively annihilated in the successive processes of $\zeta\bar{\zeta}$ recombination in bound $(\zeta\bar{\zeta})$ ζ -positronium states.

$\bar{\zeta}$ annihilation in ζ -positronium states

The frozen out antiparticles $\bar{\zeta}$ can bind at $T < I_\zeta$ with the corresponding particles ζ into positronium-like systems and annihilate. The binding is provided by the Coulomb interaction of electromagnetic charges $Z_\zeta = 2$. Since the lifetime of these positronium-like systems is much less than the timescale of their disruption by energetic photons, the direct channel of $\bar{\zeta}$ and ζ binding in $(\zeta\bar{\zeta})$, followed by a rapid annihilation, cannot be compensated by an inverse reaction of photo-ionization. That is why $\bar{\zeta}$ begins to bind with ζ and annihilates as soon as the temperature becomes less than I_ζ . The decrease of the $\bar{\zeta}$ abundance due to the $\zeta\bar{\zeta}$ recombination is governed by the equation

$$\frac{dr_{\bar{\zeta}}}{dt} = -r_{\zeta\bar{\zeta}} \cdot s \cdot \langle\sigma v\rangle, \quad (70)$$

where s is the entropy density and

$$\langle\sigma v\rangle = \left(\frac{16\pi}{3^{3/2}}\right) \cdot \frac{\bar{\alpha}}{T^{1/2} \cdot m_\zeta^{3/2}}.$$

Here $\bar{\alpha} = Z_\zeta^2 \alpha$.

In the analysis of various recombination processes, we can use the interpolation formula for the recombination cross section deduced in [3, 10, 13]:

$$\sigma_r = \left(\frac{2\pi}{3^{3/2}}\right) \cdot \frac{\bar{\alpha}^3}{T \cdot I_1} \cdot \log\left(\frac{I_1}{T}\right) \quad (71)$$

and the recombination rate given by [3, 10]

$$\langle\sigma v\rangle = \left(\frac{2\pi}{3^{5/2}}\right) \cdot \frac{\bar{\alpha}^3}{T \cdot I_1} \cdot \log\left(\frac{I_1}{T}\right) \cdot \frac{k_{in}}{M}. \quad (72)$$

Here $k_{in} = \sqrt{2TM}$, $I_1 \approx \bar{\alpha}^2 M/2$ is the ionization potential, and M is the reduced mass for a pair of recombining particles. The constant $\bar{\alpha}$ for recombining particles with charges Z_1 and Z_2 is related to the fine structure constant α via $\bar{\alpha} = Z_1 Z_2 \alpha$. The approximation Eq. (72)

followed from the known result for electron-proton recombination

$$\sigma_{rec} = \sigma_r = \sum_i \frac{8\pi}{3^{3/2}} \bar{\alpha}^3 \frac{e^4}{Mv^2 i^3} \frac{1}{(Mv^2/2 + I_i)}, \quad (73)$$

where v is the velocity of the particles. I_i is the ionization potential ($I_i = I_1/i^2$). The index i runs from one to infinity.

To sum approximately over i , it was noted in [3, 13], that $\sigma_r \propto 1/i$ for $I_i \gg Mv^2/2 = T_{eff}$, while at $I_i < T_{eff}$, the cross section $\sigma_i \propto 1/i^3$ falls down rapidly.

Using the formalism of Appendix 1, we can rewrite Eq. (70) as

$$\frac{dr_{\bar{\zeta}}}{dx} = f_{1\bar{\zeta}} \langle \sigma v \rangle r_{\bar{\zeta}} (r_{\bar{\zeta}} + \kappa), \quad (74)$$

where $x = T/I_{\zeta}$, the asymmetry $\kappa = r_{\zeta} - r_{\bar{\zeta}} = 4 \cdot 10^{-12} f_{\zeta}/S_2$ is given by Eq. (61) and

$$f_{1\bar{\zeta}} = \sqrt{\frac{\pi g_s^2}{45 g_e}} m_{Pl} I_{\zeta} \approx m_{Pl} I_{\zeta}.$$

The concentration of the remaining $\bar{\zeta}$ is given by Eq. (56) of Appendix 1

$$r_{\bar{\zeta}} = \frac{\kappa \cdot r_{f\bar{\zeta}}}{(\kappa + r_{f\bar{\zeta}}) \exp(\kappa J_{\zeta}) - r_{f\bar{\zeta}}}, \quad (75)$$

where $r_{f\bar{\zeta}}$ is given by Eq. (67) and

$$\begin{aligned} J_{\zeta} &= \int_0^{x_{f\bar{\zeta}}} f_{1\bar{\zeta}} \langle \sigma v \rangle dx = \\ &= m_{Pl} I_{\zeta} 4\pi \left(\frac{2}{3^{3/2}} \right) \cdot \frac{\bar{\alpha}^2}{I_{\zeta} \cdot m_{\zeta}} \cdot 2 \cdot x_{f\bar{\zeta}}^{1/2} \approx 0.8 \cdot 10^{15} / S_2. \end{aligned} \quad (76)$$

In the evaluation of Eq. (76) we took into account that the decrease of $\bar{\zeta}$ starts at $T \sim I_{\zeta}$, so that $x_{f\bar{\zeta}} \sim 1$. At $S_2 < 57 f_{\zeta}^{1/2}$, the abundance of $\bar{\zeta}$ is suppressed exponentially.

Indeed, one has $\kappa J_{\zeta} \approx 3200 f_{\zeta}/S_2^2$ in the exponent of Eq. (75). Similar to the case of the AC-leptons [13], it differs significantly from the situation revealed in [10] for the tera-positrons in Glashow's sinister model [9]. Though in both cases a decrease of antiparticles due to the formation of positronium like systems is induced by electromagnetic interaction and the factor in the exponent is determined by the square of the fine structure constant α , in the case of ζ -leptons, this factor is enhanced by $Z_{\zeta}^4 = 16$ times due to the Z_{ζ}^4 dependence of $\bar{\alpha}^2$. It results in a much wider mass interval for ζ -leptons, in which the primordial pair abundance is exponentially suppressed.

At S_2 close to 1, the condition $r_{f\bar{\zeta}} \ll \kappa$ in the solution Eq. (75) provides the approximate solution

$$r_{\bar{\zeta}} = r_{f\bar{\zeta}} \cdot \exp(-\kappa J_{\zeta}) \approx 10^{-14} S_2^3 \exp(-3200 f_{\zeta}/S_2^2).$$

For $S_2 > 5$, the condition $r_{f\bar{\zeta}} \gg \kappa$ is valid. Therefore the solution Eq. (75) has the form

$$r_{\bar{\zeta}} \approx \frac{\kappa}{\exp(\kappa J_{\zeta}) - 1}, \quad (77)$$

which gives for $S_2 < 57 f_{\zeta}^{1/2}$

$$r_{\bar{\zeta}} = \kappa \cdot \exp(-\kappa J_{\zeta}) \approx \left(\frac{2 \cdot 10^{-12}}{S_2} \right) \exp(-3200 f_{\zeta}/S_2^2).$$

At large $S_2 > 57 f_{\zeta}^{1/2}$, the approximate solution is given by

$$r_{\bar{\zeta}} \approx \frac{1}{J_{\zeta}} - \frac{\kappa}{2} \approx 1.25 \cdot 10^{-15} S_2 - 2 \cdot 10^{-12} f_{\zeta}/S_2.$$

In the result, the residual amount of $\bar{\zeta}$ remains at $S_2 > 57 f_{\zeta}^{1/2}$ enormously high, being for $S_2 > 70 f_{\zeta}^{1/2}$ larger than the AC-lepton excess. This effect grows with $S_2 > 70 f_{\zeta}^{1/2}$ as $\propto S_2^2$.

The general expression for the ζ abundance r_{ζ} after the ζ -positronium annihilation has the form (see Eq. (56) of Appendix 1)

$$r_{\zeta} = \frac{\kappa \cdot r_{\zeta f}}{r_{\zeta f} - (r_{\zeta f} - \kappa) \exp(-\kappa J_{\zeta})},$$

where J_{ζ} is given by Eq. (76) and $r_{\zeta f}$ is given by Eq. (67). With the account for $r_{\zeta f} > \kappa$, for all S_2 , one obtains

$$r_{\zeta} = \frac{\kappa}{1 - \exp(-\kappa J_{\zeta})}. \quad (78)$$

This gives $r_{\zeta} \approx 1/J_{\zeta} + \kappa/2 \approx 3 \cdot 10^{-16} S_2 + 2 \cdot 10^{-12} f_{\zeta}/S_2$ for large S_2 , and κ for $S_2 < 57 f_{\zeta}^{1/2}$.

-
- [1] D. Fargion et al., JETP Lett. **69**, 434 (1999); arXiv:astro-ph-9903086; K.M.Belotsky, M.Yu.Khlopov and K.I.Shibaev, Gravitation and Cosmology **6** Supplement, 140 (2000); K.M.Belotsky, D. Fargion, M.Yu. Khlopov and R.Konoplich, “May Heavy neutrinos solve underground and cosmic ray puzzles?,” arXiv:hep-ph/0411093, to appear in Phys.Atom.Nucl.; K.M.Belotsky, D.Fargion, M.Yu.Khlopov, R.Konoplich, and K.I.Shibaev, Gravitation and Cosmology **11**, 16 (2005) and references therein.

- [2] K.M. Belotsky, M.Yu. Khlopov, S.V. Legonkov and K.I. Shibaev, *Gravitation and Cosmology* **11**, 27 (2005); astro-ph/0504621.
- [3] K.M.Belotsky, D.Fargion, M.Yu.Khlopov, R.Konoplich, M.G.Ryskin and K.I.Shibaev, *Gravitation and Cosmology* **11**, 3 (2005).
- [4] M. Maltoni et al., *Phys.Lett. B* **476**, 107 (2000); V.A. Ilyin et al., *Phys.Lett. B* **503**, 126 (2001); V.A. Novikov et al., *Phys.Lett. B* **529**, 111 (2002); *JETP Lett.* **76**, 119 (2002).
- [5] M.Yu. Khlopov, *JETP Lett.* **83**, 1 (2006) [*Pisma Zh. Eksp. Teor. Fiz.* **83**, 3 (2006)]; arXiv:astro-ph/0511796
- [6] K. Belotsky, M. Khlopov and K. Shibaev, “Stable matter of 4th generation: Hidden in the Universe and close to detection?,” arXiv:astro-ph/0602261.
- [7] K. Belotsky, M. Khlopov and K. Shibaev, *Gravitation and Cosmology* **12**, 1 (2006); arXiv:astro-ph/0604518.
- [8] M. Y. Khlopov, “New symmetries in microphysics, new stable forms of matter around us,” arXiv:astro-ph/0607048.
- [9] S. L. Glashow, “A sinister extension of the standard model to $SU(3) \times SU(2) \times SU(2) \times U(1)$,” arXiv:hep-ph/0504287.
- [10] D. Fargion and M. Khlopov, “Tera-leptons shadows over sinister Universe,” arXiv:hep-ph/0507087.
- [11] C. A. Stephan, “Almost-commutative geometries beyond the standard model,” arXiv:hep-th/0509213.
- [12] A. Connes, *Noncommutative Geometry*, Academic Press, London and San Diego, 1994.
- [13] D. Fargion, M. Khlopov and C. A. Stephan, *Class. Quantum Grav.* **23**, 7305 (2006); arXiv:astro-ph/0511789.
- [14] M. Y. Khlopov and C. A. Stephan, “Composite dark matter with invisible light from almost-commutative geometry,” arXiv:astro-ph/0603187.
- [15] F. Sannino and K. Tuominen, *Phys. Rev. D* **71**, 051901 (2005); arXiv:hep-ph/0405209.
- [16] D. K. Hong, S. D. H. Hsu and F. Sannino, *Phys. Lett. B* **597**, 89 (2004); arXiv:hep-ph/0406200.
- [17] D. D. Dietrich, F. Sannino and K. Tuominen, *Phys. Rev. D* **72**, 055001 (2005); arXiv:hep-ph/0505059.
- [18] D. D. Dietrich, F. Sannino and K. Tuominen, “Light composite Higgs and precision electroweak measurements on the Z resonance: An update,” arXiv:hep-ph/0510217. To appear in

PRD.

- [19] S. B. Gudnason, C. Kouvaris and F. Sannino, Phys. Rev. D **73**, 115003 (2006); arXiv:hep-ph/0603014.
- [20] S. B. Gudnason, C. Kouvaris and F. Sannino, Phys. Rev. D **74**, 095008 (2006); arXiv:hep-ph/0608055.
- [21] V. K. Dubrovich and M. Y. Khlopov, JETP Lett. **77**, 335 (2003) [Pisma Zh. Eksp. Teor. Fiz. **77**, 403 (2003)]; arXiv:astro-ph/0206138; V. K. Dubrovich, D. Fargion and M. Y. Khlopov, Astropart. Phys. **22**, 183 (2004); arXiv:hep-ph/0312105; V. K. Dubrovich, D. Fargion and M. Y. Khlopov, Nucl. Phys. Proc. Suppl. **136**, 362 (2004).
- [22] S. Dimopoulos et al., Phys. Rev. **D41**, 2388 (1990).
- [23] C.B. Dover, T.K. Gaisser and G. Steigman, Phys. Rev. Lett. **42**, 1117 (1979); S. Wolfram, Phys. Lett. **B82**, 65 (1979); G.D. Starkman et al., Phys. Rev. **D41**, 3594 (1990); D. Javorsek et al., Phys. Rev. Lett. **87**, 231804 (2001); S. Mitra, Phys. Rev. **D70**, 103517 (2004); arXiv:astro-ph/0408341.
- [24] B.D. Wandelt et al., “Self-interacting dark matter,” arXiv:astro-ph/0006344; P.C. McGuire and P.J. Steinhardt, “Cracking open the window for strongly interacting massive particles as the halo dark matter,” arXiv:astro-ph/0105567; G. Zaharijas and G.R. Farrar, Phys. Rev. **D72**, 083502 (2005); arXiv:astro-ph/0406531.
- [25] R. Foadi, M. T. Frandsen, T. A. Rytlov and F. Sannino, “Minimal Walking Technicolor: Set Up for Collider Physics,” arXiv:0706.1696 [hep-ph].
- [26] C. Kouvaris, Phys. Rev. D **76**, 015011 (2007); arXiv:hep-ph/0703266.
- [27] N. J. Evans, S. D. H. Hsu and M. Schwetz, “Lattice tests of supersymmetric Yang-Mills theory?,” arXiv:hep-th/9707260.
- [28] K. Kainulainen, K. Tuominen and J. Virkajarvi, “The WIMP of a minimal technicolor theory,” arXiv:hep-ph/0612247.
- [29] D. S. Akerib *et al.* [CDMS Collaboration], Phys. Rev. Lett. **93**, 211301 (2004); arXiv:astro-ph/0405033.
- [30] C. Kouvaris, “WIMP Annihilation and Cooling of Neutron Stars,” arXiv:0708.2362 [astro-ph].
- [31] J. A. Harvey and M. S. Turner, Phys. Rev. D **42**, 3344 (1990).
- [32] S. M. Barr, R. S. Chivukula and E. Farhi, Phys. Lett. B **241**, 387 (1990).
- [33] S. Y. Khlebnikov and M. E. Shaposhnikov, Phys. Lett. B **387**, 817 (1996);

arXiv:hep-ph/9607386.

- [34] R. S. Chivukula and T. P. Walker, Nucl. Phys. B **329**, 445 (1990).
- [35] M. L. Burns and R. V. E. Lovelace, Astrophys.J. **202**, 87 (1982).
- [36] F. A. Agaronian and V. V. Vardanian (1985), Preprint eFI-827-54-85-YEREVAN.
- [37] M. Pospelov, Phys. Rev. Lett. **98**, 231301 (2007); arXiv:hep-ph/0605215 (2006).
- [38] M. Kaplinghat and A. Rajaraman, Phys. Rev. D **74**, 103004 (2006); arXiv:astro-ph/0606209 (2006).
- [39] K. Kohri and F. Takayama, Phys. Rev. D **76**, 063507 (2007); arXiv:hep-ph/0605243 (2006).
- [40] K. Hamaguchi, T. Hatsuda, M. Kamimura, Y. Kino and T. T. Yanagida, Phys. Lett. B **650**, 268 (2007); arXiv:hep-ph/0702274.
- [41] M. Kawasaki, K. Kohri and T. Moroi, Phys. Lett. B **649**, 436 (2007); arXiv:hep-ph/0703122.
- [42] K. Jedamzik, “The cosmic 6Li and 7Li problems and BBN with long-lived charged massive particles,” arXiv:0707.2070 [astro-ph].
- [43] M. Khlopov, Y. Levitan, E. Sedelnikov and I. Sobol, Phys.Atom.Nucl. **57**, 1393 (1994).
- [44] E. V. Sedelnikov, S. S. Filippov and M. Y. Khlopov, Phys. Atom. Nucl. **58**, 235 (1995).
- [45] M. Y. Khlopov, *Cosmoparticle physics* World Scientific, Singapore, 1999.
- [46] M. Kawasaki, K. Kohri and T. Moroi, Phys. Lett. **B625**, 7 (2005); arXiv:astro-ph/0402490.
- [47] M. Kawasaki, K. Kohri and T. Moroi, Phys. Rev. **D71**, 083502 (2005); arXiv:astro-ph/0408426.
- [48] K. Kohri, T. Moroi and A. Yotsuyanagi, Phys. Rev. **D73**, 123511 (2006); arXiv:hep-ph/0507245.
- [49] K. Jedamzik, Phys. Rev. D **74**, 103509 (2006); arXiv:hep-ph/0604251.
- [50] P. Mueller, Phys.Rev.Lett. **92**, 22501 2004; arXiv:nucl-ex/0302025.
- [51] D. McCammon et al., Nucl. Instr. Meth. **A370**, 266 (1996); D. McCammon et al., Astrophys. J. **576**, 188 (2002); arXiv:astro-ph/0205012.
- [52] C. B. Winkelman, Y. M. Bunkov, and H. Godfrin, Gravitation and Cosmology **11**, 87 (2005)
- [53] K. Belotsky, Yu. Bunkov, H. Godfrin, M. Khlopov and R. Konoplich, “He-3 experimentum crucis for dark matter puzzles,” arXiv:astro-ph/0606350.
- [54] Y. Zeldovich and I. Novikov, *Struktura i Evolyutsiya Vselennoi (in Russian), Structure and Evolution of the Universe*, Nauka, Moscow (Russian original), The University of Chicago Press, 1975 (Russian original), 1983.

- [55] R. J. Scherrer and M. S. Turner, Phys. Rev. **D33**, 1585 (1986).
- [56] A. D. Dolgov, Phys. Rept. **370**, 333 (2002); arXiv:hep-ph/0202122.